

Comparative Study of Phaco-Trabeculectomy and Small Incision Cataract Surgery Trabeculectomy in 50 Cataract Patients

Introduction

Cataracts and glaucoma often coexist, necessitating a surgical approach that addresses both conditions simultaneously. Phaco-trabeculectomy and SICS trabeculectomy are two surgical techniques that have gained prominence in managing cataracts with coexisting glaucoma.

Phaco-trabeculectomy combines phacoemulsification and trabeculectomy, offering the advantage of minimal invasiveness and quicker recovery. In contrast, SICS trabeculectomy, while effective, may be associated with a longer recovery time and higher complication rates.

This study aims to provide a comparative analysis of the two surgical techniques in terms of intraocular pressure control, visual acuity improvement, surgical time, and postoperative complications, offering insights into the optimal surgical choice for cataract patients with glaucoma.

Methodology

Study Design

A prospective, randomized study was conducted at DVVPF's Medical College & Hospital between December 2023 and November 2024. The study included 50 cataract patients diagnosed with coexisting glaucoma.

Patient Selection

- **Inclusion Criteria:**
 - Patients aged 50-80 years with confirmed cataract and glaucoma.
 - No previous ocular surgery.
 - Informed consent obtained.
- **Exclusion Criteria:**
 - Secondary glaucoma.
 - Advanced diabetic retinopathy.
 - Other ocular comorbidities affecting vision.

Surgical Procedures

- **Phaco-Trabeculectomy Group (n=25):**
 - Standard phacoemulsification with foldable IOL implantation.

- 34 ○ Trabeculectomy performed with mitomycin-C application.
- 35 ○ Sutures applied as required.
- 36 • **SICS-Trabeculectomy Group (n=25):**
- 37 ○ Manual small incision cataract surgery with rigid IOL implantation.
- 38 ○ Trabeculectomy with mitomycin-C.
- 39 ○ Suturing according to standard protocol.

40 Outcome Measures

- 41 1. **Intraocular Pressure (IOP):** Measured preoperatively, and postoperatively at 1
- 42 week, 1 month, 3 months, and 6 months using Goldmann applanation tonometry.
- 43 2. **Visual Acuity (VA):** Assessed using Snellen's chart at the same intervals.
- 44 3. **Surgical Time:** Recorded from incision to the closure of the surgical site.
- 45 4. **Complications:** Documented intraoperative and postoperative complications,
- 46 including hyphema, shallow anterior chamber, and infection.

47 Statistical Analysis

48 paired t-test was used to compare preoperative and postoperative IOP and VA within each
 49 group. An independent t-test was utilized to compare outcomes between the two groups. A p-
 50 value < 0.05 was considered statistically significant.

51

52 Results

53 Demographics and Baseline Characteristics

Parameter	Phaco-Trab Group (n=25)	SICS-Trab Group (n=25)	p-value
Age (mean ± SD)	67.4 ± 8.2	68.1 ± 7.5	0.721
Male/Female Ratio	14/11	13/12	0.793
Preoperative IOP (mmHg)	23.8 ± 2.4	24.1 ± 2.1	0.598
Preoperative VA (logMAR)	0.54 ± 0.12	0.56 ± 0.11	0.455

54

55 Intraocular Pressure (IOP) Reduction

Time Point	Phaco-Trab Group (mmHg)	SICS-Trab Group (mmHg)	p-value
Preoperative	23.8 ± 2.4	24.1 ± 2.1	0.598
1 Week Post-op	15.3 ± 1.8	16.5 ± 2.0	0.032*
1 Month Post-op	14.9 ± 1.7	16.2 ± 1.8	0.017*
3 Months Post-op	14.5 ± 1.6	15.9 ± 1.7	0.014*
6 Months Post-op	14.3 ± 1.5	15.7 ± 1.6	0.011*

56 *Statistically significant at $p < 0.05$.

57 **Visual Acuity Improvement**

Time Point	Phaco-Trab Group (logMAR)	SICS-Trab Group (logMAR)	p-value
Preoperative	0.54 ± 0.12	0.56 ± 0.11	0.455
1 Week Post-op	0.38 ± 0.10	0.42 ± 0.12	0.094
1 Month Post-op	0.34 ± 0.09	0.39 ± 0.11	0.047*
3 Months Post-op	0.32 ± 0.08	0.37 ± 0.10	0.039*
6 Months Post-op	0.31 ± 0.07	0.36 ± 0.09	0.034*

58 *Statistically significant at $p < 0.05$.

59 **Surgical Time**

Parameter	Phaco-Trab Group (minutes)	SICS-Trab Group (minutes)	p-value
Surgical Time	45.2 ± 6.4	59.1 ± 7.3	$<0.001^*$

60 *Statistically significant at $p < 0.05$.

61 **Complications**

Complication Type	Phaco-Trab Group (n=25)	SICS-Trab Group (n=25)
Hyphema	2	4
Shallow Anterior Chamber	1	3
Endophthalmitis	0	1
Choroidal Detachment	1	2
Total	4	10

62

63

64

65

66 **Discussion**

67 This study demonstrates the comparative efficacy of phaco-trabeculectomy and SICS
68 trabeculectomy in managing cataracts with coexisting glaucoma. The phaco-trabeculectomy
69 group showed superior outcomes in terms of IOP reduction, visual acuity improvement, and
70 fewer complications.

71 **Intraocular Pressure Control:** Phaco-trabeculectomy consistently achieved lower
72 postoperative IOP levels, likely due to the precise control offered by phacoemulsification and

73 the effective filtration provided by trabeculectomy. This finding aligns with previous studies
74 highlighting the effectiveness of combined procedures in managing intraocular pressure in
75 glaucoma patients.

76 **Visual Acuity Improvement:** Patients in the phaco-trabeculectomy group experienced a
77 more significant improvement in visual acuity, reflecting the minimal invasiveness and
78 superior IOL technology associated with phacoemulsification. The use of foldable IOLs in
79 phaco-trabeculectomy offers better optical quality and visual outcomes.

80 **Surgical Time:** The shorter surgical time in the phaco-trabeculectomy group is noteworthy,
81 reflecting the efficiency of modern phacoemulsification techniques. Reduced surgical time
82 correlates with less intraoperative stress and faster patient recovery.

83 **Complications:** The phaco-trabeculectomy group had fewer complications, suggesting a
84 more favorable safety profile. The higher complication rates in the SICS-trabeculectomy
85 group may be attributed to the manual nature of the procedure and the larger incision
86 required, which increases the risk of postoperative issues.

87 **Limitations:** This study's limitations include a relatively small sample size and a single-
88 center design, which may limit the generalizability of the findings. Further multicenter
89 studies with larger cohorts are needed to validate these results.

90

91 Conclusion

92 Phaco-trabeculectomy is superior to SICS trabeculectomy in managing cataract patients with
93 coexisting glaucoma, offering better intraocular pressure control, visual acuity outcomes, and
94 fewer complications. These findings support the adoption of phaco-trabeculectomy as a
95 preferred surgical approach in this patient population.

96

97 References

- 98 1. **Lichter PR, Musch DC, Gillespie BW, et al.** (2001). Interim clinical outcomes in
99 the Collaborative Initial Glaucoma Treatment Study comparing initial treatment
100 randomized to medications or surgery. *Ophthalmology*, 108(11), 1943-1953.
- 101 2. **Singh K, Shrivastava A.** (2009). Intraocular pressure fluctuations: how do they affect
102 the optic nerve? *Current Opinion in Ophthalmology*, 20(2), 90-94.
- 103 3. **Zhao Y, Chen X, Wang X.** (2017). Comparison of visual outcomes and
104 complications after phacoemulsification versus manual small incision cataract surgery
105 in highly myopic eyes. *BMC Ophthalmology*, 17(1), 65.
- 106 4. **Francis BA, Singh K, Lin SC, et al.** (2011). Novel glaucoma procedures: a report by
107 the American Academy of Ophthalmology. *Ophthalmology*, 118(7), 1466-1480.
- 108 5. **Tham CC, Kwong YY, Leung DY, et al.** (2009). Phacoemulsification versus manual
109 small-incision cataract surgery for coexisting cataract and glaucoma: a prospective,
110 randomized controlled trial. *American Journal of Ophthalmology*, 148(3), 368-375.

UNDER PEER REVIEW IN IJAR