

Observational clinical practice study of a single Adant injection in rhizarthrosis

by Jana Publication & Research

Submission date: 13-Feb-2025 12:14PM (UTC+0700)

Submission ID: 2576797428

File name: IJAR-50249.docx (126.66K)

Word count: 3934

Character count: 23195

Observational clinical practice study of a single Adant injection in rhizarthrosis

Background: Rhizarthrosis, or osteoarthritis (OA) of the first carpometacarpal joint, is currently considered a common pathology, associated with aging, affecting approximately 7% of men and 15% of women over the age of 50. Despite its high prevalence, only a small number of patients seek medical attention.

Adequate conservative treatment can alleviate symptoms and even reduce the need for surgery in up to 70% of cases. The possibilities of conservative treatment include intra-articular (IA) injection of corticosteroids and hyaluronic acid (HA). The efficacy of IA HA has been proven especially in knee but the data in rhizarthrosis are scarce.

Most HAs had been registered as medical devices worldwide and the EU Medical Devices Regulation (MDR 745/2017) requires a continuous post-market follow-up to ensure the safety and performance of these products. In compliance with the MDR, this work aimed to evaluate the efficacy and the impact on quality of life, of a marketed HA, in a cohort of patients with rhizarthrosis under real conditions in clinical practice.

Methods: Observational, post-marketing, retrospective, follow-up study. Between January 2020/June 2022, patients were treated in the Rheumatology Dpt. of Hospital General Universitario de Elche, Spain, with a single injection of HA (Adant®, Meiji Pharma Spain) and followed for 6 months. Pre/post Visual Analogue Scale (VAS) for pain, and functional questions (key, grip and button) were used for efficacy assessment. Patients' data were pseudonymized and included in a database for further analysis. The chi² test was used, for qualitative variables, and the T or Mann Whitney tests for quantitative ones. The pre/post comparison of the VAS was made with the t

test for paired samples.²⁰ The study was approved by the Ethics Committee of the hospital.

Results: Twenty patients¹ with a mean age of 61 years, 80% women, were studied. An 80% had bilateral rhizarthrosis, 70% had Kellgren-Lawrence grade III-IV (moderate-severe) and 65% had other chronic medical condition (e.g., hand OA, osteoporosis, diabetes, etc.). The 80% of the patients had received 2 previous treatments with HA injections. The volume administered varied from 1 to 2 ml (55%-45%).

The mean absolute¹ change from baseline in VAS pain score over 6 months was -5.95, a reduction of 77% ($p<0.001$). A 35% of the patients had an improvement $\geq 80\%$ and the others between 70 and 79%. Regarding functional capacities, the 80% of the patients achieved complete recovery. No significant statistical correlations were observed between baseline characteristics, the number of prior injections, or the volume administered, and the degree of improvement in pain or functional outcomes.

All patients were satisfied with the treatment. There were no adverse events recorded.

Conclusion:¹ This study suggests that viscosupplementation using Adant® is an effective and well tolerated therapeutic option in managing pain and improving function of rhizarthrosis with an excellent safety profile.

Keywords: osteoarthritis, real clinical practice, hyaluronic acid, injections, intra-articular, pain, rhizarthrosis

Introduction

Rhizarthrosis, or OA⁷ of the first carpometacarpal joint, is currently considered a common condition associated with aging, affecting approximately 7%¹⁶ of men and 15% of women over the age of 50.(1) Despite its high prevalence, only a small number of patients seek medical attention.(2)

Patients typically present with severe pain and functional impairment, and it has been shown that appropriate conservative treatment can alleviate symptoms and even

reduce the need for surgery by up to 70% in some cases.(3) HA injections are included among the conservative treatment options.(4)

Viscosupplementation with IA HA injections for OA began in the late 20th century, aiming to restore altered synovial fluid.(5) ¹³ Several studies have confirmed that HA interacts with inflammation mediators, reduces apoptosis in cartilage, stimulates chondrocyte growth, and enhances extracellular matrix protein synthesis.(5) Currently, it is a well-known and widely used treatment, particularly in knee OA, and is recognized by most scientific societies.(6–10)

In the case of rizartrrosis, studies have shown functional improvements ²⁶ in patients with moderate to severe stages of the disease, with 3 HA injections spaced 7 days apart.(11) When comparing HA use with corticosteroids, although both treatments show symptom reduction in the early weeks, after six months, results are better with HA(12), also leading ¹⁷ to a reduction in the use of Non-Steroidal Anti-Inflammatory Drugs (NSAIDs).(13)

Adant[®] is a biotechnological HA product manufactured by Meiji Pharma Spain (MPS), authorized in Europe in 1996 as a medical device for treating OA in various synovial joints.(14) Typically, it is administered intraarticularly ¹⁴ once a week for 3-5 consecutive weeks.(15) In patients with knee OA, repeated administration of Adant[®] has been shown to have a cumulative "carry over" effect, extending symptom improvement for up to one year after the last injection.(16)

Available experience with Adant[®] demonstrates its efficacy in ¹⁵ reducing symptoms and improving the quality of life in patients with rhizartrrosis.(17,18) These results, along with an excellent safety profile, make the risk/benefit ratio highly favorable.(14)

Objective

The objective is to follow a cohort of patients treated with Adant[®] for rhizartrrosis under real-world conditions. This study is part of the post-marketing surveillance of Adant[®], in accordance with the EU MDR 745/2017.(19) This new regulation, which came into force in 2023, highlights the need to follow-up medical devices throughout their lifecycle to monitor their safety and efficacy.

Material and methods

Ethics

¹⁰ The study was authorized by the research ethics committee of the *Hospital General Universitario de Elche* ³ and the waiver of informed consent was approved (approval code PI 79/2022). The study adhered to the Declaration of Helsinki. To ensure patient's privacy, data were pseudonymized.

Study design

An observational, post-marketing, ³ cross-sectional, and retrospective study in a cohort of patients with rhizarthrosis treated with Adant®.

Inclusion criteria consisted of adult patients with rhizarthrosis, confirmed radiologically, at least grade II according to the Kellgren-Lawrence classification, treated with Adant® at the *Hospital General Universitario de Elche* ¹ between January 1st, 2020, and June 30th, 2022, ⁷ with a minimum follow-up period of 6 months under routine clinical practice. Additionally, patients were required to have a VAS pain score ≥ 4 , before the treatment. The following information were obtained from patient's medical history: age, sex, severity of the infiltrated joint assessed by radiography (X-ray) according to the Kellgren-Lawrence classification, VAS pain score (baseline and after 6 months), chronic medical conditions, concomitant medication for rhizarthrosis, pathologies detected after HA injection (accidents, falls, or surgeries), infiltrated joint (left/right), administered volume (ml), previous HA injections and quality of life and satisfaction assessment questionnaire at 6 months after treatment (annex I). Patients for whom insufficient information was available were excluded from the analysis.

¹ The above information was included in a database for subsequent statistical analysis. The data were pseudonymized and identified by a code, ensuring that no information could identify the patients. Once the data has been analyzed, a statistical report was prepared, which served as the basis for the subsequent publication of the results.

Outcomes measures

The VAS was used to measure pain. Pain measured by VAS consisted of a discrete scale from 0 (no pain) to 10 (maximum pain).

The quality of life and satisfaction assessment questionnaire was conducted as part of routine clinical practice during the patient's visit to the specialist to assess hand functionality and treatment satisfaction in patients with rhizarthrosis after six months of follow-up. The questions were completed by the physician during the interview to evaluate functional recovery in daily activities and the patient's subjective perception of treatment outcomes. While not part of a standardized tool, it included elements inspired by validated instruments such as Functional Index for Hand Osteoarthritis (FIHOA) and Michigan Hand Outcomes Questionnaire (MHQ).

Statistics

Descriptive Statistics

Qualitative variables were described using absolute frequencies and percentages. Quantitative variables that follow a normal distribution was described using mean, standard deviation (SD), minimum (Min), and maximum (Max); while those that do not follow a normal distribution was described using median, interquartile range (first quartile (Q1) – third quartile (Q3)), Min, and Max.

Analytical Statistics

Univariate comparisons between categorical variables were performed using the chi-square test and/or Fisher's exact test. For continuous variables, the shape of the distributions was analyzed using the Kolmogorov-Smirnov and Shapiro-Wilk tests. Comparisons between two unrelated means were made using the Student's t-test or the Mann-Whitney U test. In the case of analyzing more than two groups, comparisons were made using ANOVA or the Kruskal-Wallis test.

Results

Study population: sociodemographic and clinical characteristics

The study included a cohort of 20 patients with a mean age of 63.1 years (SD = 7.76). The 80% of the patients (16) were women. The severity of rhizarthrosis was classified

using Kellgren-Lawrence system, with the 55% of patients being grade III. All sociodemographic and clinical features were detailed in table 1.

Chronic medical condition

The chronic medical condition presented in the patients at the beginning of the treatment under study were registered, 13 patients (65%) had at least one comorbidity. Nine patients (45%) presented some osteoarticular pathology. Seven patients (35%) had no chronic medical condition. The main ones included hand OA and osteoporosis. (Table 2).

Previous treatment with hyaluronic acid

Regarding previous HA injections received by the patients for rhizarthrosis, 80% (16) of them had previously received a median of 2 injections. The 10% of the patients (2) received 3 injections, one patient received 4 injections, and another patient received 8 previous injections.

Treatment with hyaluronic acid in study

During the study, the patients received a HA injection for the treatment of rhizarthrosis as part of routine clinical practice. The 40% of the patients (8) received the treatment in both first carpometacarpal joints. The median injected HA volume was 1 ml. The 55% of the patients (11) received 1 ml, and the 45% (9) were injected with a volume of 2 ml.

Concomitant medication

All patients were undergoing additional treatments concomitantly with the administration of the HA injection as part of their overall therapeutic approach. Twelve patients (60%) were treated with NSAIDs, specifically etoricoxib. The concomitant medication during the study were summarized in the table 3.

Efficacy assessment

The mean baseline VAS score was 7.70 (SD = 1.17), and the mean final VAS score at 6 months was 1.75 (SD = 1.07) ($p < 0.001$), (figure 1).

All patients showed improvements in the VAS, with the smallest reduction being 28.6%. Reductions of at least 30% were considered clinically relevant. Only one patient did not achieve this improvement. This patient was a 62-year-old woman with rhizarthrosis Kellgren-Lawrence grade II, who had received two prior treatment with HA. She presented with a baseline VAS score of 7, and from a functional perspective, she did not have problems with key handling, buttoning, or grip closure.

A reduction $\geq 80\%$ in the VAS was considered a high improvement. This level of improvement was achieved by 35% of the patients (7) in the study. Thirteen patients showed an improvement in the VAS between 70% and 79%. The mean pain reduction was 77% in the study population ($p < 0.001$).

Regarding the recovery of joint functionality at 6 months, three questions were assessed (key handling, grip the fist, and buttoning), and it was recorded whether the patient was able to perform each activities or not. Sixteen patients (80%) demonstrated adequate functionality in all three activities at the 6-month follow-up, while 20% (4) showed partial improvement in joint functionality, achieving improvement in at least one of the three activities. (Table 4)

All the patients were satisfied with the HA treatment.

No predictive factors of response were identified among the variables measured in the study (basal characteristics, number of previous injections and volume administered) and the pain or function improvements of rhizarthrosis.

Safety

During the study, no adverse events were recorded. No patient developed any other significant health issues after receiving the HA injection, including accidents, falls, or surgeries.

24

Discussion

The results of this study provide data on the management of rhizarthrosis with a single HA injection in real-world clinical practice. Currently, there is limited evidence compared to its use in knee OA.(6–10) Our findings suggest that HA injection into the

first carpometacarpal joint could represent ¹² an effective treatment for improving pain and functionality in patients with rhizarthrosis, offering a less invasive option before considering surgical treatments.

Many studies have demonstrated the beneficial effects of HA in knee OA, with ²⁷ improvements in pain and joint function. However, extrapolating these results to rhizarthrosis is not straightforward due to biomechanical differences and variations in joint load.

Fuchs ²⁵ et al., conducted a prospective, controlled, randomized study assessed the efficacy and tolerability of IA HA and ¹ triamcinolone acetonide in 56 patients with OA of the carpometacarpal joint of the thumb over 26 weeks. Patients received three injections. Results showed that triamcinolone acetonide provided quicker pain relief at 2-3 weeks, while HA showed a slight superiority at week 26 and had significantly better lateral pinch power. ¹² Heyworth et al. in a double-blind controlled trial, included 60 patients with basal joint OA. Patients were randomized to receive two injections of HA, ¹ one saline injection followed by a corticosteroid injection, or two saline injections. No statistically significant between-group differences in pain were observed, but significant improvements in pain compared to baseline at weeks 12 and 26 were seen in the HA group. ²⁰ In the study of Figen et al., ³² 33 women with bilateral thumb base OA were included and received single injection of HA in one hand, and saline in the other hand. ⁶ Statistically significant improvements were detected in function ($p = 0.001$), VAS pain ($p = 0.002$), and pinch strength ($p = 0.004$) at the 24th week in the HA group. However, only VAS pain scores decreased temporarily in control hands at the 6th week ($p = 0.02$). ²¹ Bahadir et al conducted a ⁴ randomized, open-label, evaluator-blinded clinical study including 40 women with stage II or III trapeziometacarpal joint OA. The steroid group ($n = 20$) received one injection of 20 mg triamcinolone acetonide once and the HA group ($n = 20$) received three injections of 5 mg HA at 1-week intervals. Pain level decreased significantly over 12 months for the steroid group and over 6 months for the HA group. Hand function improved in both groups but it was only significant in the steroid group. ⁸ ²² Velasco et al. in a prospective, single-arm, multicenter, open-label study with a 6-month follow-up period included 35 patients with rhizarthrosis treated with a single HA injection.

¹ The least-squares mean change from baseline in VAS pain score over 6 months was –2.00, a reduction of 27.8% (p<0.001).(23)

These studies present promising results, though with limitations, including small sample sizes and methodological heterogeneity. Our findings align ¹⁴ with the results of these studies, further supporting the potential efficacy of HA injections in the management of rhizarthrosis. Additionally, our data provide ²⁹ valuable insights into the real-world, contributing to the evidence and complementing existing knowledge.

¹ The patients included in this study reported a high level of satisfaction with HA injection, highlighting ¹¹ significant improvements in pain and quality of life. Furthermore, no adverse events were recorded, suggesting an excellent safety profile for this treatment. This aspect is particularly important in long-term treatments for chronic conditions such as OA. Moreover, a favorable safety profile enhances both healthcare professionals' and patients' confidence in the treatment, contributing to better adherence and satisfaction.

Comparisons with clinical trials should be made with caution, as both the design and methodology of observational real-world studies differ substantially from those of randomized clinical trials. In this context, clinical trials typically limit or prohibit the use of prior HA injections, which ¹² may not reflect the real-world clinical practice. In our study, the majority of the patients included had previously received IA HA treatments.

On the other hand, our study has some limitations inherent to its retrospective nature and clinical practice design. Patients were managed at the discretion of the physician based on clinical practice, rather than according to a pre-established protocol. Additionally, the data were retrospectively collected from the available medical records, following the routine practice at the participating site. Knowledge of clinical practice is crucial for understanding how treatments are applied in real-world and how healthcare professionals make decisions based on experiences and outcomes. In the context of HA injections, this knowledge enables physicians to identify which patients may benefit the most, treatments to their specific needs, and improve health outcomes while ensuring adherence to best clinical practices based on scientific evidence.

Conclusion

Six months after a single HA injection, all patients had benefited from treatment, either due to decreased pain, improved functional abilities, or both.

No predictive factors of response were identified among the variables measured in the study and the pain or function improvements of HA injection.

HA did not appear to have a negative effect on significant health issues after receiving the HA injection, including accidents, falls, or surgeries.

The safety profile of HA injections in clinical practice was favorable and similar to that previously described.

In conclusion, ⁷our findings suggest that HA injection could be an useful treatment in the management of rhizarthrosis, helping to improve pain and functionality with an excellent safety profile. ³⁰However, further research is required to establish its long-term impact and its positioning within clinical guidelines.

⁵Declarations

Ethics approval and consent to participate

The study was authorized by the research ethics committee of the Hospital General Universitario de Elche ³ and the waiver of informed consent was approved (approval code PI 79/2022).

⁵Consent for publication

Not applicable.

Availability of data and materials

The datasets used and analyzed during the current study are available from the authors on reasonable request.

Competing interests

MP. Coronel Granada and D. Acosta-Rubio³ are employees of Meiji Pharma Spain.

Other authors do not have COI.

Funding

This study was funded by Meiji Pharma Spain (MPS).

Authors' contribution

Concept and design of the study: CGMP, NBF, NPJR

Collection of data: NBF, NPJR

Interpretation of the data: NBF, NPJR, ARD, CGMP

Drafting of the manuscript: ARD, CGMP

Critical revision and final approval: NBF, NPJR

Acknowledgements

The statistical analysis and report were conducted by D. J. J. Granizo Martínez (Granadatos SL).

List of abbreviations

HA - hyaluronic acid

IA - intra-articular

IQR - Interquartile Range

Max - maximum

MDR - medical devices regulation

Min - minimum

MPS – Meiji Pharma Spain

NSAIDs - Non-steroidal anti-inflammatory drugs

OA – osteoarthritis

19

Q1 – first quartile

Q3 – third quartile

SD - standard deviation

VAS - Visual Analog Scale

Tables and figures

Table 1. Sociodemographic and clinical characteristics

Variable	N=20
Age (years), mean (SD)	63.1 (7.76)
Female sex, n (%)	16 (80)
Affected joint, n (%)	
Left	5 (25)
Right	7 (35)
Both	8 (40)
Kellgren-Lawrence grade, n (%)	
II	6 (30)
III	11 (55)
IV	3 (15)
Chronic medical condition, n (%)	13 (65)

Number of prior injections, median (IQR)	2 (2-8)
Injected volumen (ml), median (IQR)	1 (1-2)

Table 2. Chronic medical condition

Chronic medical condition	N (%)
Hand osteoarthritis*	3 (15)
Osteoporosis*	3 (15)
Diabetes	2 (10)
Chondromalacia*	1 (5)
Gonarthrosis*	1 (5)
Hypercholesterolemia	1 (5)
Hypertension	1 (5)
Palindromic rheumatism*	1 (5)
Rheumatoid arthritis*	1 (5)

* Articular pathology

Table 3. Concomitant medication


Concomitant medication	N (%)
NSAIDs	12 (60)
Paracetamol	4 (20)
Nutraceutical	3 (15)
Etanercept	1 (5)

Multiple options possible per patient.

Table 4. Functional assessment

Adequate function	Yes, n (%)	No, n (%)
Key	18 (90)	2 (10)
Grip	19 (95)	1 (5)
Button	17 (85)	3 (15)

Figure 1. Mean of VAS pain score



References

1. Haara MM, Heliövaara M, Kröger H, Arokoski JPA, Manninen P, Kärkkäinen A, et al. Osteoarthritis in the Carpometacarpal Joint of the Thumb: Prevalence and Associations with Disability and Mortality. *JBJS* [Internet]. 2004;86(7). Available from: https://journals.lww.com/jbjsjournal/fulltext/2004/07000/osteoarthritis_in_the_carpometacarpal_joint_of_the.13.aspx
2. Wilkens SC, Meghpara MM, Ring D, Coert JH, Jupiter JB, Chen NC. Trapeziometacarpal arthrosis. *JBJS Rev.* 2019;7(1):e8.
3. Berggren Jane Lindstrand Goran Nylander Bo Povlsen Magnus AJD. Reduction in the need for operation after conservative treatment of osteoarthritis of the first carpometacarpal joint: a seven year prospective study. *Scand J Plast Reconstr Surg Hand Surg.* 2001;35(4):415–7.
4. Fuggle N, Bere N, Bruyère O, Rosa MM, Prieto Yerro MC, Dennison E, et al. Management of hand osteoarthritis: from an US evidence-based medicine guideline to a European patient-centric approach. *Aging Clin Exp Res.* 2022;34(9):1985–95.
5. Altman RD, Manjoo A, Fierlinger A, Niazi F, Nicholls M. The mechanism of action for hyaluronic acid treatment in the osteoarthritic knee: a systematic review. *BMC Musculoskelet Disord.* 2015;16:1–10.
6. Conrozier T, Monfort J, Chevalier X, Raman R, Richette P, Diraçoğlu D, et al. EUROVISCO recommendations for optimizing the clinical results of viscosupplementation in osteoarthritis. *Cartilage.* 2020;11(1):47–59.

7. Trojian TH, Concoff AL, Joy SM, Hatzenbuehler JR, Saulsberry WJ, Coleman CI. AMSSM scientific statement concerning viscosupplementation injections for knee osteoarthritis: importance for individual patient outcomes. *Br J Sports Med.* 2016;50(2):84–92.
8. Bannuru RR, Osani MC, Vaysbrot EE, Arden NK, Bennell K, Bierma-Zeinstra SMA, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis Cartilage.* 2019;27(11):1578–89.
9. Phillips M, Bhandari M, Grant J, Bedi A, Trojian T, Johnson A, et al. A systematic review of current clinical practice guidelines on intra-articular hyaluronic acid, corticosteroid, and platelet-rich plasma injection for knee osteoarthritis: an international perspective. *Orthop J Sports Med.* 2021;9(8):23259671211030270.
10. Bruyère O, Honvo G, Veronese N, Arden NK, Branco J, Curtis EM, et al. An updated algorithm recommendation for the management of knee osteoarthritis from the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO). In: *Seminars in arthritis and rheumatism.* Elsevier; 2019. p. 337–50.
11. Mandl LA, Hotchkiss RN, Adler RS, Lyman S, Daluiski A, Wolfe SW, et al. Injectable hyaluronan for the treatment of carpometacarpal osteoarthritis: open label pilot trial. *Curr Med Res Opin.* 2009;25(9):2103–8.
12. Fuchs S, Mönikes R, Wohlmeiner A, Heyse T. Intra-articular hyaluronic acid compared with corticoid injections for the treatment of rhizarthrosis. *Osteoarthritis Cartilage* [Internet]. 2006;14(1):82–8. Available from: <https://www.sciencedirect.com/science/article/pii/S1063458405002001>
13. Frizziero A, Maffulli N, Masiero S, Frizziero L. Six-months pain relief and functional recovery after intra-articular injections with hyaluronic acid (mw 500–730 kDa) in trapeziometacarpal osteoarthritis. *Muscles Ligaments Tendons J.* 2014;4(2):256.
14. ADANT® (Hyaluronic Acid). Clinical Evaluation Report 2021. Data on file.
15. Product Information. Adant® (Sodium Hyaluronate for Intra-articular Administration). Data on file.
16. Navarro-Sarabia F, Coronel P, Collantes E, Navarro FJ, de la Serna AR, Naranjo A, et al. A 40-month multicentre, randomised placebo-controlled study to assess the efficacy and carry-over effect of repeated intra-articular injections of hyaluronic acid in knee osteoarthritis: the AMELIA project. *Ann Rheum Dis.* 2011;70(11):1957–62.
17. de Leon Garcia FJ, Fernandez MAR, Rodriguez RZ, Gonzalez CTY. Tratamiento con ácido hialurónico en la rizartrrosis del pulgar. *Medicina de Rehabilitación.* 2003;16(1):8–12.
18. Conca Roig A. Tratamiento de la artrosis trapecio-metacarpiana con ácido hialurónico bajo control radiológico. *Revista Española de Reumatología* 25;5. 1998 May;
19. Regulation (EU) 2017/745 of the European Parliament and of the Council of 5 April 2017 on Medical Devices, Amending Directive 2001/83/EC, Regulation (EC) No 178/2002 and Regulation (EC) No 1223/2009 and Repealing Council Directives 90/385/EEC and

93/42/EEC [Internet]. 2017 p. 1–175. Available from:
<http://data.europa.eu/eli/reg/2017/745/oj>

20. Heyworth BE, Lee JH, Kim PD, Lipton CB, Strauch RJ, Rosenwasser MP. Hylan Versus Corticosteroid Versus Placebo for Treatment of Basal Joint Arthritis: A Prospective, Randomized, Double-Blinded Clinical Trial. *J Hand Surg Am* [Internet]. 2008;33(1):40–8. Available from:
<https://www.sciencedirect.com/science/article/pii/S0363502307009124>
21. Figen Ayhan F, Üstün N. The evaluation of efficacy and tolerability of Hylan G-F 20 in bilateral thumb base osteoarthritis: 6 months follow-up. *Clin Rheumatol* [Internet]. 2009;28(5):535–41. Available from: <https://doi.org/10.1007/s10067-008-1080-0>
22. Bahadır C, Onal B, Dayan VY, Güreş N. Comparison of therapeutic effects of sodium hyaluronate and corticosteroid injections on trapeziometacarpal joint osteoarthritis. *Clin Rheumatol* [Internet]. 2009;28(5):529–33. Available from:
<https://doi.org/10.1007/s10067-008-1079-6>
23. Velasco E, Ribera MV, Pi J. Single-arm open-label study of Durolane (NASHA nonanimal hyaluronic acid) for the treatment of osteoarthritis of the thumb. *Open Access Rheumatol* [Internet]. 2017 Mar 27;9(null):61–6. Available from:
<https://www.tandfonline.com/doi/abs/10.2147/OARRR.S128675>

Observational clinical practice study of a single Adant injection in rhizarthrosis

ORIGINALITY REPORT

27%	24%	23%	4%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	www.dovepress.com Internet Source	7%
2	www.mdpi.com Internet Source	3%
3	Amparo Santamaría Torroba, María Pilar Acin Lazaro, Elena Gómez-Rubio, Pilar Coronel Granado. "Efficacy and Safety of One Shot of Hyaluronic Acid in Hip Osteoarthritis: Postmarketing Clinical Follow-Up for Real-World Evidence", Open Access Rheumatology: Research and Reviews, 2024 Publication	2%
4	www.researchgate.net Internet Source	1%
5	www.researchsquare.com Internet Source	1%
6	journals.sagepub.com Internet Source	1%
7	www.science.gov Internet Source	1%
8	Submitted to University of Edinburgh Student Paper	1%
9	www.accc.gov.au Internet Source	1%

10	www.revistafarmaciahospitalaria.es Internet Source	1 %
11	discovery.researcher.life Internet Source	1 %
12	www.frontiersin.org Internet Source	1 %
13	www.journalijar.com Internet Source	1 %
14	Francesco Ioppolo, Fabiana Saracino, Rosaria Sabrina Rizzo, Giampaolo Monacelli et al. "Comparison Between Extracorporeal Shock Wave Therapy and Intra-articular Hyaluronic Acid Injections in the Treatment of First Carpometacarpal Joint Osteoarthritis", <i>Annals of Rehabilitation Medicine</i> , 2018 Publication	1 %
15	"World Congress on Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (WCO-IOF-ESCEO 2023)", <i>Aging Clinical and Experimental Research</i> , 2023 Publication	<1 %
16	docksci.com Internet Source	<1 %
17	dmoco.com Internet Source	<1 %
18	docusalut.com Internet Source	<1 %
19	www.jstage.jst.go.jp Internet Source	<1 %
20	www.prolekare.cz Internet Source	<1 %

21	doczz.net Internet Source	<1 %
22	repub.eur.nl Internet Source	<1 %
23	link.springer.com Internet Source	<1 %
24	openaccess.leidenuniv.nl Internet Source	<1 %
25	"Articular Cartilage Defects of the Knee", Springer Nature, 2012 Publication	<1 %
26	aprenderly.com Internet Source	<1 %
27	chk.static.cigna.com Internet Source	<1 %
28	mfprm.org Internet Source	<1 %
29	Alejandro Alcina Navarro, Sara Gómez Valero, Mercedes Gimeno del Sol, María Pilar Coronel Granado. "Navigating the New EU Medical Devices Regulation: Retrospective Post- Market Follow-Up of Hyaluronic Acid Injections for Knee Osteoarthritis", Open Access Rheumatology: Research and Reviews, 2024 Publication	<1 %
30	Viktor Shtroblia, Pavlo Petakh, Iryna Kamyshna, Iryna Halabitska, Oleksandr Kamyshnyi. "Recent advances in the management of knee osteoarthritis: a narrative review", Frontiers in Medicine, 2025 Publication	<1 %

31

Sara Tenti, Sara Cheleschi, Nicola Mondanelli, Stefano Giannotti, Antonella Fioravanti. "New Trends in Injection-Based Therapy for Thumb-Base Osteoarthritis: Where Are We and where Are We Going?", *Frontiers in Pharmacology*, 2021

Publication

<1 %

32

www.ncbi.nlm.nih.gov

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On