

1 **“Knowledge and attitude on neonatal danger signs among the antenatal women residing**
2 **in the urban areas of Gangtok, Sikkim”**
3

UNDER PEER REVIEW IN IJAR

4 **Abstract:**

5 **Background:** A child's survival during the first month of life is highly dependent on several
6 factors since it's a very fragile and delicate period. According to WHO 47% of all under 5
7 deaths occurred in the newborn period and majority of the deaths occurred in the developing
8 country. The mother has a direct impact on her child's health, thus she has to be alert to any
9 early warning indications that could be concerning.

10
11
12 **Aim:** The aim of the study was to assess the knowledge and attitude possessed on neonatal
13 danger signs by the antenatal women.

14
15
16 **Materials and Methods:** A descriptive cross-sectional study was conducted among 237
17 antenatal women (Primigravida and multigravida) residing in the urban areas of Gangtok,
18 Sikkim. The samples were collected using Purposive sampling technique. A structured
19 knowledge questionnaire and 5-point Likert scale was administered to collect the data.

20
21
22 **Results:** The results showed that about 5% of the antenatal women had good knowledge, 77%
23 had average knowledge and 18% had poor knowledge, whereas majority, 97% of the women
24 possessed favourable attitude and only 3% had unfavourable attitude. There was a statistically
25 significant correlation ($r = 0.317$ at 0.001 level of significance) between knowledge and attitude
26 on neonatal danger signs.

27
28
29 **Conclusion:** The study concluded that there is a need to improve knowledge on neonatal
30 danger signs among the antenatal women. This can be achieved by providing education either
31 during ANC visits, PNC follow-ups or at community level. Therefore, interventional strategies
32 that stresses on strengthening maternal education should be extended.

33
34
35 **Key words:** neonatal danger signs, knowledge, attitude, antenatal women, urban areas

Introduction

“Infant mortality and life expectancy are reasonable indicators of general well-being in a society.”

The first month of life is an extremely important and fragile time in a child's existence. In addition to being a time for development and connection, it is also a time to exercise prudence. The period of first 28 days of the newborn's existence is referred to as neonatal phase. ⁽¹⁾

The transition from intrauterine to extra uterine life is a significant hurdle or a struggle for the baby after birth. WHO in 2020 reported that 2.4 million new-borns died in the first 28 days of life, accounting for over half (47%) of all fatalities among children under the age of five. Neonatal danger signs are non-specific signs of severe illness and can be a manifestation of almost any newborn disease that can be easily identified by non- clinical personnel which includes the mother. ⁽²⁾

Neonatal danger signs are clinical indicators suggesting a high risk of neonatal morbidity and mortality. Most neonatal fatalities take place in underdeveloped nations, whereby the majority of the deaths taking place in home. Even though there has been progress where the mortality rate dropped from 38.8 deaths per 1,000 live births in 2013 to 26.619 deaths in 2023, but still there remains a gap in number of fatalities across various nations ⁽³⁾

One of the health-related Sustainable Development Goals (SDG) that must be accomplished by 2030 is lowering newborn mortality. Therefore, enhancing neonatal health and lowering mortality rate plays a pivotal role in achieving the SDG. WHO estimates that half of all deaths occurred in only five developing nations namely Ethiopia, India, Nigeria, Pakistan and Democratic Republic of Congo. ⁽⁴⁾

World Health Organization (WHO) and United Nations International Children's Emergency Fund (UNICEF) have recommended neonatal danger signs, which warns that the new-borns are at high risk of sickness and death. In 2013 the World Health Organization issued strong recommendations for the assessment of particular danger signs during each postnatal visit. If any of these signs are present, the newborn should be immediately referred for further evaluation. ⁽⁵⁾

The new-born deaths were caused by number of causes that includes, limited awareness and understanding among the primary care providers especially the mothers, difficulty in reaching a conclusive diagnosis and lack of specificity in the clinical presentations of numerous neonatal morbidities. ⁽⁶⁾

The mother has an increased responsibility to recognize the danger signs because of the brief hospital stay after childbirth and the restricted time for a thorough examination of the new-born. According to some studies, most new-born deaths in the developing countries occur as a result of mothers' failure to recognize these danger signs and not seeking medical care at the earliest. ⁽⁷⁾

Neonates are often hospitalized with variety of symptoms that indicate illness. These symptoms may be present during the time of hospital stay or may develop after the baby is discharged from the hospital. Here, the aim of initial management of a neonate showing these symptoms is stabilization and preventing deterioration of the health. Neonates are more prone to show subtle signs of illness and difficulty of feeding that are sometimes the only signs present, but the illness may advance quickly ⁽⁸⁾

It is estimated that 75% of neonatal deaths could be avoided with simple low-cost identification and management tools and this is only possible if the mothers' gain knowledge regarding the above neonatal signs of danger (NSD) that enable them to make a quick and prompt decision of seeking health assistance. The World Health Organization (WHO) developed Integrated Management of New-born Illness initiative, which focused on the assessment of NSD and application of prompt timely treatment. ⁽⁹⁾

The health condition of a child directly contacts with the mother, she must be very aware of the early signs which could also possibly be a warning sign. Mother or the caregivers are the first people who can notice the early warning characteristics of illness and major changes or deviation from normal. The main aim is early recognition of the occurrence of these danger signs that would aid in predicting the need for seeking treatment of the new-born. ⁽¹⁰⁾

Early detection of neonatal danger signs of illness is a very important step towards maximizing the neonate's chances of survival. It should be noted that most of the neonates in the developing countries are either delivered at home or are discharged from the health facility too early. Intervention modalities that focus on increasing the level of prenatal education, access to antenatal and postnatal care and advocating the use of television as means for health education were pinpointed in some relevant studies. ⁽¹¹⁾

The objectives of this study were:

1. To estimate the knowledge and attitude on neonatal danger signs among the antenatal women.
2. To determine the correlation between knowledge and attitude on neonatal danger signs.
3. To determine the association between the knowledge on neonatal danger signs with demographic and obstetrics variables.
4. To determine the association between the attitude on neonatal danger signs with demographic and obstetrics variables.

Materials and method

This study is a descriptive cross-sectional study and adopted purposive sampling technique. The study setting was the urban areas of Gangtok, Sikkim where 237 samples were selected based on the inclusion and exclusion criteria. The inclusion criteria were as follows: primi and multigravida antenatal women of all the trimester, antenatal women who are willing to participate, between the age group of 19-45 years and antenatal women who understands either English, Hindi and Nepali language.

The data collection tool in the study included: Tool I: Section A: Demographic profile of the antenatal women and Section B: Obstetrics profile of the antenatal women

Demographic profile included components like age, religion, marital status, educational status of the women and husband, occupation of women and husband, total family income and type of family.

Obstetrics profile included components like age at first pregnancy, parity, history of abortion, history of still birth, number of living children, frequency of antenatal visits, plan for place of delivery, antenatal visit accompanied by spouse and birth preparedness status.

Tool II: Structured knowledge questionnaire on neonatal danger signs which included 11 components give by WHO (not being able to feed, or stopped feeding well, convulsed or fitted since birth, fast breathing, chest in drawing, high temperature or low temperature, yellow soles, movement only when stimulated, or no movement even on stimulation and signs of local infection such as umbilicus redness or draining of pus, skin boils or eyes draining pus)

Tool III: 5- point Likert scale on neonatal danger signs which included components like danger signs presented by newborn, seek medical care, time taken to visit a health facility, reasons to not seek health facility, exposure to sunlight, frequency of breastfeeding, need for education during antenatal period.

The study was conducted at the urban areas of Gangtok, Sikkim. Administrative approval from the concerned authorities and respective councillors of the area was taken. The purpose of the study was explained to all the participants after which an informed consent was taken. Patient information sheet was given to the participants where the objectives, procedure involved and their right to withdraw any moment from the study was explained. The tools were then administered and data were collected using interview technique

The data was analysed using descriptive and inferential statistics in SPSS 2024. Chi square or Fischer's exact test was computed to find out the association between knowledge and attitude with demographic and obstetrics variables. Karl Pearson's Correlation Coefficient was used to measure the correlation between knowledge and attitude.

Results

Based on the results related to demographic profile of antenatal women, majority 51% of the participants belonged to the age group of 28-37 years with 54% belonging to Hindu and all the antenatal women were married. Majority 28% of the women had secondary and above graduation qualification and 29% of their husband had senior secondary level qualifications. Majority 56% of them were homemaker while 35% of their husbands works at other firms. Approximately 35% of the participants total income of family was above 30,001 and majority 51% of them belonged to a joint family

Based on the results related to obstetrics profile of antenatal women, majority 60% of the participants were in the age group of 19-27 years when they had their first pregnancy. 57% of the women were primigravida and 43% of them were multigravida Majority 91% of them had no history of abortion and only 0.8% of them had a history of stillbirth. Majority 65% of the participants had a frequency of antenatal visit more than four and 94% of the women were accompanied by their spouses during their visits. Majority 93% of the participants claimed that it was a planned pregnancy and 64% of them had preferred government setting for their place of delivery.

Based on the results related to knowledge and attitude on neonatal danger signs; the results revealed that out of 237 participants, only 5% had good knowledge, 77% had average knowledge and 18% of them had poor knowledge on neonatal danger signs. The findings also revealed that majority of the participants 97% possessed favorable attitude and only 3% of them had unfavorable attitude on neonatal danger signs.

Based on the results related to correlation between knowledge and attitude, there was a moderate positive correlation between the knowledge and attitude since the obtained value of 'r' and 'p' was ($r = 0.317$) ($p = 0.001$).

Based on the results, there was a statistically significant association found between knowledge and demographic variables (age in years, educational status of both the women husband, occupation of antenatal women, occupation of husband and total income of family. A statistically significant association was also found between knowledge and women's age at first pregnancy whereas no significant association was found with other obstetric variables.

Based on the results, there was a statistically significant association found between attitude and the educational status of both antenatal women and husband. The results also revealed that there was no any significant association between attitude and obstetrics variables.

Table 1: Frequency and percentage distribution of demographic variables

N= 237

| Sl. No | Socio-demographic Variables | Frequency (<i>f</i>) | Percentage (%) |
|--------|---------------------------------------|---------------------------|-------------------|
| 1. | Age in years | | |
| | a. 19-27 | 96 | 41 |
| | b. 28-37 | 121 | 51 |
| | c. 38-45 | 20 | 8 |
| 2. | Religion | | |
| | a. Hindu | 127 | 54 |
| | b. Christian | 53 | 23 |
| | c. Buddhist | 51 | 21 |
| | d. Others | 6 | 2 |
| 3. | Marital status | | |
| | a. Married | 237 | 100 |
| | b. Unmarried | 0 | 0 |
| | c. Divorced/ Separated | 0 | 0 |
| | d. Widowed | 0 | 0 |
| 4. | Educational status of antenatal women | | |
| | a. No formal education | 13 | 5 |
| | b. Primary education | 32 | 14 |
| | c. Secondary | 67 | 28 |
| | d. Senior secondary | 58 | 24 |
| | e. Graduation and above | 68 | 29 |
| 5. | Educational status of husband | | |
| | a. No formal education | 17 | 7 |
| | b. Primary education | 32 | 14 |
| | c. Secondary | 61 | 26 |
| | d. Senior secondary | 69 | 29 |
| | e. Graduation and above | 58 | 24 |

| | | | |
|----|-------------------------------|------------|-----------|
| 6. | Occupation of antenatal women | | |
| | a. Homemaker | 134 | 56 |
| | b. Private employee | 39 | 17 |
| | c. Government employee | 52 | 22 |
| | d. Others | 12 | 5 |
| 7. | Occupation of husband | | |
| | a. Unemployed | 16 | 7 |
| | b. Private employee | 72 | 30 |
| | c. Government employee | 65 | 28 |
| | d. Others | 84 | 35 |
| 8. | Total income of family | | |
| | a. ≤10,000 | 24 | 10 |
| | b. 10,001- 20,000 | 65 | 27 |
| | c. 20,001-30,000 | 66 | 28 |
| | d. Above 30,001 | 82 | 35 |
| 9. | Type of family | | |
| | a. Nuclear | 97 | 41 |
| | b. Joint | 120 | 51 |
| | c. Extended | 20 | 8 |

Table 2: Frequency and percentage distribution of Obstetrics variables

N= 237

| Sl. No | Obstetrics Variables | Frequency (<i>f</i>) | Percentage (%) |
|--------|------------------------------|---------------------------|-------------------|
| 1. | Age at first pregnancy | | |
| | a. 19-27 | 141 | 60 |
| | b. 28-37 | 92 | 38 |
| | c. 38-45 | 4 | 2 |
| 2. | Number of pregnancies | | |
| | a. Primi | 134 | 57 |
| | b. Multi | 103 | 43 |
| 3. | History of abortion | | |
| | a. Yes | 22 | 9 |
| | b. No | 215 | 91 |
| 4. | History of still birth | | |
| | a. Yes | 2 | 1 |
| | b. No | 235 | 99 |
| 5. | No. of living children | | |
| | a. None | 136 | 57 |
| | b. 1 | 83 | 35 |
| | c. 2 | 17 | 7 |
| | d. 3 and above | 1 | 1 |
| 6. | Frequency of antenatal visit | | |
| | a. Less than 4 visits | 84 | 36 |
| | b. More than 4 visits | 153 | 64 |
| 7. | Plan for place of delivery | | |
| | a. Government hospital | 151 | 64 |

| | | | |
|----|---------------------------------------|------------|-----------|
| | b. Private hospital | 86 | 36 |
| | c. Home setting | 0 | 0 |
| | d. Others | 0 | 0 |
| 8. | Antenatal visit accompanied by spouse | | |
| | a. Yes | 224 | 95 |
| | b. No | 13 | 5 |
| 9. | Birth preparedness status | | |
| | a. Planned | 220 | 93 |
| | b. Unplanned | 17 | 7 |

Table 3: Frequency and percentage distribution of level of knowledge on neonatal danger signs among the antenatal women

N= 237

| Knowledge | Frequency (f) | percentage (%) | Score Range | Median | Mode | Mean | SD |
|-----------|------------------|-------------------|----------------|--------|------|------|------|
| Poor | 43 | 18 | 0-14 | | | | |
| Average | 182 | 77 | 14 | 8 | 8 | 8.10 | 2.78 |
| Good | 12 | 5 | | | | | |

Table 4: Area wise distribution of knowledge on neonatal danger signs among antenatal women.

N= 237

| Area of knowledge | Total items | Total Maximum Score | Total score obtained | Mean | Mean % |
|--|--------------------|----------------------------|-----------------------------|-------------|---------------|
| Previous knowledge and meaning on neonatal danger signs | 2 | 2 | 176 | 0.74 | 37% |
| Poor sucking or has stopped feeding | 3 | 3 | 342 | 1.44 | 48% |
| Signs of convulsions | 1 | 1 | 121 | 0.51 | 51% |
| Fast and difficult breathing | 3 | 3 | 312 | 1.31 | 44% |
| Hyperthermia | 2 | 2 | 224 | 0.94 | 47% |
| Hypothermia | 1 | 1 | 113 | 0.47 | 48% |
| Yellow skin and soles | 1 | 1 | 126 | 0.53 | 53% |
| Treatment for jaundice | 1 | 1 | 86 | 0.36 | 36% |

| | | | | | |
|---------------------------|---|---|-----|------|-----|
| Unconsciousness | 2 | 2 | 199 | 0.83 | 42% |
| Signs of infection | 2 | 2 | 211 | 0.89 | 45% |

Section III: Description of attitude on neonatal danger signs among the antenatal women

N= 237




Figure 1: Frequency percentage distribution of attitude on neonatal danger signs among the antenatal women

Table 5: Correlation between knowledge and attitude on neonatal danger signs among the antenatal women

N= 237

| Variables | Mean | SD | r value | p value |
|-----------|-------|------|---------|---------|
| Knowledge | 8.10 | 2.78 | 0.317 | 0.001* |
| Attitude | 73.70 | 5.59 | | |




Figure 2: correlation between knowledge and attitude on neonatal danger signs among the antenatal women

Table 6: Association between knowledge on neonatal danger signs with demographic variables.

N= 237

| Sl. no. | Demographic variables | Poor knowledge | Average knowledge | Good knowledge | Fischer's exact/ χ^2 value | df | p value |
|---------|---------------------------------------|----------------|-------------------|----------------|------------------------------------|----|---------------------|
| 1. | Age in years | | | | | | |
| | a. 19-27 | 23 | 69 | 4 | 8.898 | 4 | 0.048* |
| | b. 28-37 | 20 | 93 | 8 | | | |
| | c. 38-45 | 0 | 20 | 0 | | | |
| 2. | Religion | | | | | | |
| | a. Hindu | 24 | 98 | 5 | 5.231 | 6 | 0.473 ^{NS} |
| | b. Christian | 8 | 41 | 4 | | | |
| | c. Buddhist | 8 | 40 | 3 | | | |
| | d. Others | 3 | 3 | 0 | | | |
| 3. | Marital status | | | | | | |
| | a. Married | 43 | 182 | 12 | NA | NA | NA |
| | b. Unmarried | 0 | 0 | 0 | | | |
| | c. Divorced/ Separated | 0 | 0 | 0 | | | |
| | d. Widowed | 0 | 0 | 0 | | | |
| 4. | Educational status of antenatal women | | | | | | |
| | a. No formal education | 8 | 5 | 0 | 65.41 | 8 | 0.001* |
| | b. Primary education | 17 | 15 | 0 | | | |
| | c. Secondary | 11 | 55 | 1 | | | |
| | d. Senior secondary | 7 | 50 | 1 | | | |
| | e. Graduation and above | 0 | 57 | 10 | | | |
| 5. | Educational status of husband | | | | | | |
| | a. No formal education | 5 | 12 | 0 | 35.68 | 8 | 0.001* |
| | b. Primary education | 16 | 16 | 0 | | | |
| | c. Secondary | 9 | 50 | 2 | | | |

| | | | | | | | |
|-----------|-------------------------------|-----------|-----------|-----------|-------|---|---------------------|
| | d. Senior secondary | 11 | 55 | 3 | | | |
| | e. Graduation and above | 2 | 49 | 7 | | | |
| 6. | Occupation of antenatal women | | | | | | |
| | a. Homemaker | 34 | 98 | 2 | 28.93 | 6 | 0.001* |
| | b. Private | 4 | 34 | 1 | | | |
| | c. Government | 2 | 44 | 6 | | | |
| | d. Others | 3 | 6 | 3 | | | |
| 7. | Occupation of husband | | | | | | |
| | a. Unemployed | 6 | 10 | 0 | 16.78 | 6 | 0.006* |
| | b. Private | 9 | 61 | 2 | | | |
| | c. Government | 5 | 55 | 5 | | | |
| | d. Others | 23 | 56 | 5 | | | |
| 8. | Total income of family | | | | | | |
| | a. ≤10,000 | 8 | 16 | 0 | 27.41 | 6 | 0.001* |
| | b. 10,001- 20,001 | 20 | 44 | 1 | | | |
| | c. 20,001-30,000 | 10 | 55 | 1 | | | |
| | d. Above 30,001 | 5 | 67 | 10 | | | |
| 9. | Type of family | | | | | | |
| | a. Nuclear | 17 | 75 | 5 | 1.076 | 4 | 0.911 ^{NS} |
| | b. Joint | 21 | 93 | 6 | | | |
| | c. Extended | 5 | 14 | 1 | | | |

Table 7: Association between knowledge on neonatal danger signs with Obstetrics variables

N= 237

| Sl. no. | Obstetrics variables | Poor knowledge | Average knowledge | Good knowledge | Fischer's exact/ χ^2 value | df | p value |
|-----------|------------------------------|----------------|-------------------|----------------|------------------------------------|----|---------------------|
| 1. | Age at first pregnancy | | | | | | |
| | d. 19-27 | 35 | 99 | 7 | 10.85 | 4 | 0.021* |
| | e. 28-37 | 8 | 79 | 5 | | | |
| | f. 38-45 | 0 | 4 | 0 | | | |
| 2. | Number of pregnancies | | | | | | |
| | a. Primi | 20 | 106 | 8 | 2.432 | 2 | 0.314 ^{NS} |
| | b. Multi | 23 | 76 | 4 | | | |
| 3. | History of abortion | | | | | | |
| | a. Yes | 2 | 19 | 1 | 1.225 | 2 | 0.574 ^{NS} |
| | b. No | 41 | 163 | 11 | | | |
| 4. | History of still birth | | | | | | |
| | a. Yes | 0 | 2 | 0 | 0.968 | 2 | 0.995 ^{NS} |
| | b. No | 43 | 180 | 12 | | | |
| 5. | Number of living children | | | | | | |
| | a. None | 20 | 108 | 8 | 8.496 | 4 | 0.211 ^{NS} |
| | b. 1 | 20 | 59 | 4 | | | |
| | c. 2 | 2 | 15 | 0 | | | |
| | d. 3 and above | 1 | 0 | 0 | | | |
| 6. | Frequency of antenatal visit | | | | | | |
| | a. Less than 4 visits | 18 | 64 | 2 | 2.507 | 2 | 0.278 ^{NS} |
| | b. More than 4 visits | 35 | 118 | 10 | | | |
| 7. | Plan for place of delivery | | | | | | |
| | a. Government | 26 | 118 | 7 | 0.567 | 2 | 0.789 ^{NS} |
| | b. Private hospital | 17 | 64 | 5 | | | |
| | c. Home setting | 0 | 0 | 0 | | | |
| | d. Others | 0 | 0 | 0 | | | |

| | | | | | | | |
|-----------|---------------------------------------|-----------|------------|-----------|-------|---|---------------------|
| 8. | Antenatal visit accompanied by spouse | | | | | | |
| | a. Yes | 40 | 172 | 12 | 0.447 | 2 | 0.859 ^{NS} |
| | b. No | 3 | 10 | 0 | | | |
| 9. | Birth preparedness status | | | | | | |
| | a. Planned | 39 | 170 | 11 | 0.892 | 2 | 0.543 ^{NS} |
| | b. Unplanned | 4 | 12 | 1 | | | |

Table 8: Association between attitude on neonatal danger signs with demographic variables

N= 237

| Sl. no. | Demographic variables | Unfavorable attitude | Favorable attitude | Fischer's exact/χ^2value | df | p value |
|----------------|---------------------------------------|-----------------------------|---------------------------|---|-----------|---------------------|
| 1. | Age in years | | | | | |
| | a. 19-27 | 4 | 92 | 0.691 | 2 | 0.841 ^{NS} |
| | b. 28-37 | 3 | 118 | | | |
| | c. 38-45 | 0 | 20 | | | |
| 2. | Religion | | | | | |
| | a. Hindu | 5 | 122 | 5.242 | 3 | 0.098 ^{NS} |
| | b. Christian | 0 | 53 | | | |
| | c. Buddhist | 1 | 50 | | | |
| | d. Others | 1 | 5 | | | |
| 3. | Marital status | | | | | |
| | a. Married | 7 | 230 | NA | NA | NA |
| | b. Unmarried | 0 | 0 | | | |
| | c. Divorced/ Separated | 0 | 0 | | | |
| | d. Widowed | 0 | 0 | | | |
| 4. | Educational status of antenatal women | | | | | |
| | a. No formal education | 1 | 12 | 8.920 | 4 | 0.020* |
| | b. Primary education | 1 | 31 | | | |
| | c. Secondary | | | | | |

| | | | | | | |
|----|-------------------------------|----------|------------|-------|---|---------------------|
| | d. Senior secondary | 5 | 62 | | | |
| | e. Graduation and above | 0 | 58 | | | |
| | | 0 | 10 | | | |
| 5. | Educational status of husband | | | | | |
| | a. No formal education | 2 | 15 | 9.857 | 4 | 0.009* |
| | b. Primary education | 1 | 31 | | | |
| | c. Secondary | | | | | |
| | d. Senior secondary | 4 | 57 | | | |
| | e. Graduation and above | 0 | 69 | | | |
| | | 0 | 58 | | | |
| 6. | Occupation of antenatal women | | | | | |
| | a. Homemaker | 6 | 128 | 2.221 | 3 | 0.479 ^{NS} |
| | b. Private | 1 | 38 | | | |
| | c. Government | 0 | 52 | | | |
| | d. Others | 0 | 12 | | | |
| 7. | Occupation of husband | | | | | |
| | a. Unemployed | 0 | 16 | 5.059 | 3 | 0.119 ^{NS} |
| | b. Private | 5 | 67 | | | |
| | c. Government | 0 | 65 | | | |
| | d. Others | 2 | 82 | | | |
| 8. | Total income of family | | | | | |
| | a. ≤10,000 | 1 | 23 | 3.586 | 3 | 0.259 ^{NS} |
| | b. 10,001- 20,001 | 4 | 61 | | | |
| | c. 20,001-30,000 | 1 | 65 | | | |
| | d. Above 30,001 | 1 | 81 | | | |
| 9. | Type of family | | | | | |
| | a. Nuclear | 3 | 94 | 0.217 | 2 | 0.891 ^{NS} |
| | b. Joint | 4 | 116 | | | |
| | c. Extended | 0 | 20 | | | |

Table 9: Association between attitude on neonatal danger signs with Obstetrics variables

N= 237

| Sl. no. | Obstetrics variables | Unfavorable attitude | Favorable attitude | Fischer's exact/ χ^2 value | df | p value |
|---------|------------------------------|----------------------|--------------------|------------------------------------|----|---------------------|
| 1. | Age at first pregnancy | | | | | |
| | a. 19-27 | 4 | 137 | 0.734 | 2 | 0.992 ^{NS} |
| | b. 28-37 | 3 | 89 | | | |
| | c. 38-45 | 0 | 4 | | | |
| 2. | Number of pregnancies | | | | | |
| | a. Primi | 3 | 131 | 0.544 | 1 | 0.461 ^{NS} |
| | b. Multi | 4 | 99 | | | |
| 3. | History of abortion | | | | | |
| | a. Yes | 0 | 22 | 1.386 | 1 | 0.239 ^{NS} |
| | b. No | 7 | 208 | | | |
| 4. | History of still birth | | | | | |
| | a. Yes | 0 | 2 | 0.120 | 1 | 0.729 ^{NS} |
| | b. No | 7 | 228 | | | |
| 5. | Number of living children | | | | | |
| | a. None | 3 | 133 | 3.074 | 3 | 0.571 ^{NS} |
| | b. 1 | 4 | 79 | | | |
| | c. 2 | 0 | 17 | | | |
| | d. 3 and above | 0 | 1 | | | |
| 6. | Frequency of antenatal visit | | | | | |
| | a. Less than 4 visits | 2 | 82 | 0.154 | 2 | 0.695 ^{NS} |
| | b. More than 4 visits | 5 | 148 | | | |
| 7. | Plan for place of delivery | | | | | |
| | a. Government | 3 | 148 | 1.294 | 1 | 0.255 ^{NS} |
| | b. Private hospital | 4 | 82 | | | |
| | c. Home setting | 0 | 0 | | | |
| | d. Others | 0 | 0 | | | |

| | | | | | | |
|----|---------------------------------------|---|-----|-------|---|---------------------|
| 8. | Antenatal visit accompanied by spouse | 6 | 218 | 0.774 | 1 | 0.379 ^{NS} |
| | a. Yes | 1 | 12 | | | |
| | b. No | | | | | |
| 9. | Birth preparedness status | 6 | 214 | 0.438 | 1 | 0.508 ^{NS} |
| | a. Planned | 1 | 16 | | | |
| | b. Unplanned | | | | | |

Discussion in relation to knowledge on neonatal danger signs.

The findings of this study revealed that 76.8% of the participants had average knowledge, 18.1% had poor knowledge and only 5.1% had good knowledge.

These findings are supported by a study conducted at Dhulikhel, Nepal by **Ratneworee Prajapati** and **Sujata Madhikarmi** in 2016 where 55.2% of the participants' possessed moderate knowledge, 9.6% of them had poor knowledge and 35.2% of the respondents had inadequate knowledge on new-born danger signs. ⁽¹²⁾

Also, according to a study conducted at Ethiopia by **Mulugeta W, Tarikua Afework** and **Prem Kumar** in 2019, the findings showed that 82.90% were not knowledgeable and only 17.01% of the respondents were knowledgeable on the identification of new-born danger. ⁽¹³⁾

However, the findings are inconsistent with the study conducted at Jalandhar Punjab by **Balbir Singh et al.** in 2021 where 49.6% of the women had good knowledge, 38% had poor knowledge and 12.4% had zero knowledge where they couldn't identify even one neonatal danger sign. The disparity may be due to the differences in the demographic profile or operational definition in their study. ⁽¹⁴⁾

Discussion in relation to attitude on neonatal danger signs.

The findings of the present study shows that 97% of the antenatal women had a favorable attitude and only 3% of them possessed unfavorable attitude on neonatal danger signs. These findings are similar with the study conducted at Dehradun, Uttarakhand by **Reena Thakur, Rajesh Kumar, Laxmi Kumar and Sanchita Pugazhendi** in 2017 which showed majority of the respondents (61%) had moderate attitude and 39% of them had favorable attitude on neonatal danger signs. ⁽¹⁵⁾

Discussion in relation to association between knowledge on neonatal danger signs with demographic and obstetrics variables

In the present study, 51.1% of the participants belonged to the age group of 28-37 years, 40.5% between 19-27 and 8.4% between 38-45 years. Similar findings were reported by a study

conducted at Ethiopia by **Guta A, Seema A, Amsalu B and Sintayehu Y** in the year 2020, where majority of the participants (62.8%) belonged to the middle age group of 25-34 years, 23.2% between 18-24 years and 14% \geq 35 years. The present study showed that the antenatal women's age was statistically significant to their knowledge on neonatal danger signs. Antenatal women who were between the age group of 28-37 had higher level of knowledge than the women belonging to other age groups. ⁽¹⁶⁾

Discussion in relation to association between attitude on neonatal danger signs with demographic and obstetrics variables

The findings of the present study show a statistically significant association between attitude possessed on neonatal danger sign and educational status of both the antenatal women and her husband. The findings are supported by a study conducted by **Reena Thakur, et al.** in 2017 where they found statistically significant association between attitude score of the participants and level of education ($p = 0.001$). Hence it can be interpreted statistically that the mothers who have higher education were also having more positive attitude regarding neonatal danger signs. ⁽¹⁵⁾

Conclusion:

The present study concluded that, the antenatal women had an average knowledge on neonatal danger signs with a favorable attitude towards it. Even though majority of the antenatal women possessed average knowledge and notably favorable attitude towards it, still there remains a need to educate the antenatal women and strengthen interventional strategies that improve the knowledge of the antenatal women. There was a statistically significant moderate positive correlation between knowledge and attitude on neonatal danger signs which highlights the crucial interplay between knowledge and attitude.

According to the findings, there is a need to foster deeper understanding to enhance early recognition and prompt management of neonatal danger signs, ultimately contributing to improve maternal and child health outcomes. This can be achieved by stressing counselling sessions to address any misconceptions or concerns regarding neonatal health.

Acknowledgment

I express my profound gratitude whole heartedly to Ms. Roshmi Tamang, MSc. Nursing, Department of Obstetrics and Gynecological Nursing, Sikkim Manipal College of Nursing for her major contribution in the study.

I extend my sincere and heartfelt gratitude to co-guide Mrs. Sonam Zangmu Sherpa, Associate Professor, Department of Obstetrics and Gynecological Nursing, Sikkim Manipal College of Nursing for constant support, masterly guidance, valuable advices and suggestions from beginning till the end of the study.

I gratefully extend my sincere gratitude to Prof. Champa Sharma, Principal, Sikkim Manipal College of Nursing for providing the opportunity to conduct the study and providing various learning opportunities throughout the study.

I also take this opportunity to thank all the faculty members of Sikkim Manipal College of Nursing for their timely advice, keen support and motivation that helped me in the successful completion of the study

A word of sincere and heartfelt thanks to all the respective councillors of the urban areas of Gangtok, Sikkim for facilitating the necessary administrative permission to conduct this study.

I owe my deep sense of gratitude to all the validators for their constructive viewpoints on the tools selected for the study. Their valuable and constructive suggestions proved to be worthwhile.

My sincere and heartfelt thanks to Mr. Tara Prasad Dhakal, Managing Editor, Sangrila Times, Mrs. Pema Bhutia, PGT English and Mrs. Ambika Adhikari, PGT Hindi for translation of tools, patient information sheet, consent form to English, Nepali and Hindi language.

A word of thanks to the statistician Prof. Naveen Kumar, Sri Guru Ram Das University of Health Sciences, Punjab for his valuable input for the statistical analysis.

I am also sincerely grateful to my family, especially my beloved mother Mrs. Mamita Tamang and uncle Mr. Dewan Rai for their unwavering support and understanding during this journey. Their encouragement and belief in me have been the driving force to complete this study.

My special thanks to all the participants who generously shared their valuable time and participated in my study, without whom this research would not have been possible.

Prof. Binita Khati
HOD, OBG Nursing
Sikkim Manipal College of Nursing

References:

1. Pathirana J, Muñoz FM, Abbing-Karahagopian V, Bhat N, Harris T, Kapoor A, et al. Neonatal death: Case definition & guidelines for data collection, analysis, and presentation of immunization safety data. *Vaccine*. 2016 Dec 1; 34(49):6027–37.
2. World Health Organization, 2021 <https://www.who.int/news-room/fact-sheets/detail/levels-and-trends-in-child-mortality-report-2021>
3. World Health Organization, 2019, monitoring health for the SDGs, <https://www.who.int/publications/i/item/9789241565707>
4. Yosef, T., Nigussie, T., & Asefa, A. (2020). Knowledge of Neonatal Danger Signs and Its Associated Factors among Mothers Attending Child Vaccination Centers at Sheko District in Southwest Ethiopia. *International journal of pediatrics*, 2020, 40919
5. World Health Organization, postnatal care for mother and newborn <https://www.who.int/docs/default-source/mca-documents/nbh/brief-postnatal-care-for-mothers-and-newborns-highlights-from-the-who-2013-guidelines.pdf>
6. Guta, A., Sema, A., Amsalu, B., & Sintayehu, Y. Knowledge of Neonatal Danger Signs and Associated Factors among Mothers of <6 Months Old Child in Dire Dawa, Ethiopia: A Community Based Cross-Sectional Study. *International journal of women's health*. 2020; 12: 539–548.
7. Datta Parul, Textbook of Paediatric nursing, 3rd edition, New Delhi, India, Jaypee Brothers;2014.
8. Mohammed Ahmed NSO, Al-Gamar EA. Mother's knowledge and attitude regarding recognition of neonatal signs of danger. 2022; 22(1):27-35.
9. Alebel, A., Wagnew, F., Petrucka, P., Tesema, et al. (2020). Neonatal mortality in the neonatal intensive care unit of Debre Markos referral hospital, Northwest Ethiopia: a prospective cohort study. *BMC pediatrics*, 20(1), 72.
10. Ali Rehan Syed, BhuttaZulfiqar. Newborn Health in Developing Countries Hutchison's Paediatrics. 2nd edition; Jaypee Brothers Medical Publishers; 2012 January.
11. Degefa N, Diriba K, Girma T, Kebede A, Senbeto A, Eshetu E, et al. Knowledge about Neonatal Danger Signs and Associated Factors among Mothers Attending Immunization Clinic at Arba Minch General Hospital, Southern Ethiopia: A Cross-Sectional Study. *BioMed Research International*. 2019, August, 6; 2019.
12. Prajapati R, Madhikarmi S. Knowledge Regarding Newborn Danger Signs among Antenatal Mothers Attending Out Patient Department in Dhulikhel Hospital. *International Journal of Health Sciences & Research [Internet]*. 2016; 6(6):268
13. Gebre MW. Knowledge, Attitude and Practice on Identification of Newborn Danger Signs and Associated Factors Among Mothers attending Postnatal Clinics at Dessie Referral Hospital, North East Ethiopia. 2021, January 21; 4(4): 191-200.

14. Singh B, Chawla G, Kaur M, Singh J, Singh J, Sonakshi., et al. Knowledge about Danger Signs in Neonates and Health Seeking Behaviour amongst Mothers Attending Tertiary Healthcare Centre in Punjab, India. *Journal of clinical and diagnostic research*. 2021 November;15(11).
15. Thakur R, Sharma RK, Kumar L, Pugazhendi S. Neonatal Danger Signs: Attitude and Practice of Post-Natal Mothers. *Journal of Nursing & Care*. 2017 May, 19; 06(03).
16. Guta, A., Sema, A., Amsalu, B., & Sintayehu, Y. Knowledge of Neonatal Danger Signs and Associated Factors among Mothers of <6 Months Old Child in Dire Dawa, Ethiopia: A Community Based Cross-Sectional Study. *International journal of women's health*. 2020; 12: 539–548.

UNDER PEER REVIEW IN IJHR