

Atrogenic Gastric Perforation in an 18-Month-Old Infant: A Case Report

Abstract: We report the case of an 18-month-old Moroccan infant admitted for suspected foreign body ingestion, who developed an iatrogenic gastric perforation. Bronchoscopy, performed without evidence of a foreign body, was complicated by severe bronchospasm requiring intubation, mechanical ventilation, and prolonged sedation. During the ICU stay, the placement of a nasogastric tube led to gastric mucosal micro-erosions, promoting the formation of a stress ulcer and its subsequent perforation. Three days post-intubation, abdominal distension with peritoneal signs prompted a CT scan, confirming massive pneumoperitoneum. Emergency laparotomy with gastric suturing and peritoneal lavage was performed but was unsuccessful due to a fatal hypovolemic shock.

Keywords: Iatrogenic gastric perforation, infant, nasogastric tube, laparotomy, hypovolemic shock.

Introduction: Iatrogenic gastric perforation is a rare but serious complication that can be life-threatening if not managed promptly. We present the case of an infant who developed gastric perforation secondary to an intervention, highlighting clinical, radiological, and therapeutic aspects.

Methodology: This is a retrospective case report conducted in October 2024 at the Pediatric Intensive Care Unit of Abderrahim Harouchi Mother-Child Hospital, Ibn Rochd University Hospital, Casablanca. The study is based on a detailed analysis of the medical records of an 18-month-old infant admitted for suspected foreign body ingestion, who subsequently developed an iatrogenic gastric perforation.

Clinical data were comprehensively collected from the patient's medical records, including medical history, clinical and paraclinical findings, as well as diagnostic and therapeutic interventions performed during hospitalization. The patient's clinical course was chronologically reviewed to identify factors contributing to this complication.

Results: An 18-month-old Moroccan infant, with no significant medical history, was initially admitted for suspected foreign body ingestion. The patient presented with persistent cough and respiratory discomfort. Bronchoscopy was performed but revealed no foreign body. This procedure was complicated by severe bronchospasm requiring intubation, mechanical ventilation, and prolonged sedation for 48 hours.

During this period, a nasogastric tube was inserted for gastric decompression due to mechanical ventilation. However, gastric mucosal micro-erosions induced by the tube, combined with stress ulceration, led to gastric perforation.

39 Three days post-intubation, the patient developed marked abdominal distension with
40 worsening general condition. The abdomen was tympanic, and peritonitis was strongly
41 suspected. An abdominal CT scan confirmed a massive pneumoperitoneum.

42



43

44 **Figure 1:** Thoraco-abdominal CT scan showing massive pneumoperitoneum, indicative of
45 gastric perforation.

46 Before transfer to the operating room, the patient experienced cardiac arrest, which was
47 reversed following intensive resuscitation. Emergency laparotomy revealed an antral gastric
48 perforation associated with massive pneumoperitoneum and purulent intra-abdominal fluid. A
49 single-layer gastric suture was performed, along with extensive peritoneal lavage and drain
50 placement.



51

52

53 **Figure 2:** Laparotomy for pneumoperitoneum due to gastric perforation in an infant.

54 Despite surgical management and post-operative resuscitation efforts, the patient suffered
55 another cardiac arrest in the recovery room, which was irreversible. Death was declared
56 shortly after, directly resulting from complications related to massive pneumoperitoneum and
57 hypovolemic shock.

58

59 **Discussion:** Iatrogenic gastric perforation is a rare but potentially fatal complication in
60 infants. In our case, the 18-month-old infant developed gastric perforation secondary to an
61 initial intervention for suspected foreign body ingestion, complicated by nasogastric tube
62 placement and prolonged mechanical ventilation. This iatrogenic mechanism has been
63 described in the literature, with several contributing factors, including mechanical trauma
64 from repeated device insertion (1,2).

65 Nasogastric tube insertion is a common procedure in pediatric intensive care; however, it
66 carries risks such as gastric mucosal micro-erosions, which, in the presence of a stress ulcer,
67 can progress to perforation (3). Prolonged mechanical ventilation further contributes to gastric
68 distension and increased intra-abdominal pressure, facilitating perforation (4). The rapid
69 clinical deterioration of our patient, marked by severe abdominal distension and peritoneal
70 signs, underscores the importance of early imaging diagnosis, particularly CT scans, to
71 confirm massive pneumoperitoneum (5).

72 Emergency surgical management is crucial in such cases. The laparotomy performed in our
73 case, involving gastric perforation suturing and peritoneal lavage, is the recommended
74 therapeutic approach to limit peritoneal contamination (6). Despite this intervention,
75 multiorgan failure and hypovolemic shock due to generalized peritoneal infection led to the

76 patient's death. These complications have also been reported in other studies, highlighting the
77 high mortality rate in cases of delayed management (7).

78 This case underscores the importance of close monitoring and early intervention in suspected
79 iatrogenic gastric perforation. Prevention includes careful handling during invasive device
80 insertion and regular position verification to reduce microtrauma risks, particularly in infants
81 with fragile gastric walls.

82

83 **Conclusion:** Iatrogenic gastric perforation in infants remains a diagnostic and therapeutic
84 challenge requiring a multidisciplinary approach. This case report highlights the critical
85 importance of early recognition and urgent surgical intervention to improve outcomes.

86

87 **References:**

- 88 1. Yao HHI, Tuck MV, McNally C, Smith M, Usatoff V. Gastric rupture following
89 nasopharyngeal catheter oxygen delivery—a report of two cases. *Anaesth Intensive*
90 *Care*. 2015;43(3):244-248.
- 91 2. Miller JS, Itani KM, Oza MD, Wall MJ. Gastric rupture with tension
92 pneumoperitoneum: a complication of difficult endotracheal intubation. *Ann Emerg*
93 *Med*. 1997;30(5):343-346.
- 94 3. Reinhold H, Hilderbrand-Mathieu M, Rucquoi M, Bollaert A. Gastric rupture
95 complicating oxygen therapy by nasal tube. *Acta Anaesthesiol Belg*. 1973;24:241-255.
- 96 4. Barichello AW, Pimblett T, Dyck FJ, McFadden D. Rupture of the stomach following
97 oxygen therapy by nasal catheter. *Can Med Assoc J*. 1968;98(11):855-858.
- 98 5. Pue CA, Pacht ER. Complications of fiberoptic bronchoscopy at a university hospital.
99 *Chest*. 1995;107(2):430-432.
- 100 6. Fazlalizadeh H, Adimi P, Kiani A, Malekmohammad M, Jabardarjani HR,
101 Soltaninejad F, et al. Evaluation of bronchoscopy complications in a tertiary health
102 care center. *Tanaffos*. 2014;13(2):48-50.
- 103 7. Onwuka EA, Saadai P, Boomer LA, Nwomeh BC. Nonoperative management of
104 esophageal perforations in the newborn. *J Surg Res*. 2016;205(1):102-107.
- 105 8. Al-Qahtani SM, Al-Shamrani MM, Al-Harbi AM. Iatrogenic gastric perforation in
106 pediatric patients: A case series and review of the literature. *J Pediatr Surg*.
107 2019;54(6):1158-1163.
- 108 9. Lin C, Yang X, Peng C, Zhang H. Stress ulcer and gastric perforation in critically ill
109 pediatric patients: Risk factors and clinical outcomes. *World J Gastroenterol*.
110 2021;27(19):2245-2253.
- 111 10. Oh A, Gulati G, Muthuvel S, Langham MR. Gastric perforation in infants: Etiology,
112 clinical presentation, and outcomes. *J Pediatr Gastroenterol Nutr*. 2017;64(4):e101-
113 e106.

114

115

116