


# Jana Publication & Research

## Mastoid Process: Morphometric Parameters with Correlation to Side and Gender

 28

 BioTech

 Institut Seni Indonesia Surakarta

---

### Document Details

Submission ID

trn:oid::1:3167734334

Submission Date

Feb 27, 2025, 5:10 PM GMT+7

Download Date

Feb 27, 2025, 5:32 PM GMT+7

File Name

IJAR-50428.docx

File Size

1.6 MB

13 Pages

1,675 Words

8,804 Characters

# 58% Overall Similarity





The combined total of all matches, including overlapping sources, for each database.

## Filtered from the Report




- ▶ Bibliography
- ▶ Quoted Text

---

## Match Groups

-  **33 Not Cited or Quoted 31%**  
Matches with neither in-text citation nor quotation marks
-  **14 Missing Quotations 26%**  
Matches that are still very similar to source material
-  **0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
-  **0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

## Top Sources

- 57%  Internet sources
- 27%  Publications
- 6%  Submitted works (Student Papers)

### Match Groups

- **33 Not Cited or Quoted 31%**  
Matches with neither in-text citation nor quotation marks
- **14 Missing Quotations 26%**  
Matches that are still very similar to source material
- **0 Missing Citation 0%**  
Matches that have quotation marks, but no in-text citation
- **0 Cited and Quoted 0%**  
Matches with in-text citation present, but no quotation marks

### Top Sources

- 57% Internet sources
- 27% Publications
- 6% Submitted works (Student Papers)

### Top Sources

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Internet	www.sciencepub.net	30%
2	Internet	www.ncbi.nlm.nih.gov	11%
3	Internet	archive.org	8%
4	Internet	www.researchgate.net	4%
5	Internet	iafmonline.in	1%
6	Internet	rcastoragev2.blob.core.windows.net	1%
7	Internet	www.spandidos-publications.com	<1%
8	Internet	www.ijfmt.com	<1%
9	Publication	Yumi Hoshioka, Suguru Torimitsu, Yohsuke Makino, Daisuke Yajima et al. "Sex es...	<1%
10	Publication	Jigyasa Passey, Suniti Pandey, Rahul Singh, Shailendra Singh. "Morphometric Stu...	<1%

## Mastoid Process: Morphometric Parameters with Correlation to Side and Gender

**3** ABSTRACT The mastoid part is the posterior region of the temporal bone and projects down as the conical mastoid process. It is larger in adult males than in females. Sternocleidomastoid, splenius capitis and longissimus capitis muscles are all attached to its lateral surface. The posterior belly of digastric muscle is attached to a deep mastoid notch on its medial aspect. The occipital artery runs in a shallow occipital groove which lies medial to the mastoid notch. The study was conducted in Department of Anatomy, Pt. B.D. Sharma PGIMS, Rohtak, In year 23-24 on 120 dried human skulls out of which 80 were of males and 40 were of females. The study on the parameters of the mastoid process is important in the determination of sex for forensic purposes and anthropologists. It was concluded in our study that the mean mastoid parameters were more in male skulls than female skulls.

**5**

**2**

**2**

KEYWORDS Mastoid process, temporal bone

3 INTRODUCTION The mastoid part is the posterior region of the temporal bone and projects

down as the conical mastoid process. It is larger in adult males than in females. Sternocleidomastoid, splenius capitis and longissimus capitis muscles are all attached to its lateral surface. The posterior belly of digastric muscle is attached to a deep mastoid notch on its medial aspect. The occipital artery runs in a shallow occipital groove which lies medial to the mastoid notch. (Standring S.2005) For

2 many anthropologists, while excavating skeletal remains or in cases of unforeseen disasters, identification of gender is the preliminary task. A major role in the gender identification of skeletal remains may be played by morphometric osteological criteria and lays the foundation for full identification.

5 MATERIALS AND METHODS The study was conducted in Department of Anatomy, Pt. B.D. Sharma PGIMS, Rohtak, In year 23-24 on 120 dried human skulls out of which 80 were of males and 40 were of females. Skulls with broken temporal bone were excluded from the study. Mastoid process of both right and left sides were studied. Following morphometric parameters were measured using vernier caliper.

1 A. **Mastoid length:** straight distance from mastoidale to the upper rim of root of zygomatic process of temporal bone.

B. **Mastoid breadth:** the straight distance from posterior end of incisura mastoidea (digastric notch) to the nearest point of posterior border of external auditory meatus.

1 **Asterion– Mastoidale length (AST-MS):** the straight distance between asterion and mastoidale (both right and left sides). (Saadia A.2016)

**Asterion – Porion length (AST-PO):** the straight distance between asterion and porion (both right and left sides). (Saadia A.2016)

**Porion – Mastoidale length (PO-MS):** the straight distance between porion and mastoidale (right and left side). (Saadia A.2016)

4 **Porion (PO):** superior point of external auditory meatus.<sup>2</sup> **Mastoidale (MS):** most inferior point of the mastoid process.<sup>2</sup> **Asterion (AST):** the point where the parietal, temporal and occipital bones meets.

(Saadia A.2016)



8 *Figure: 1 Measurement of mastoid length*



Figure:

Figure: 2 *Measurement of mastoid breadth*





Figure: 3 Measurement of Asterion– Mastoidale length (AST-MS)



Figure: 4 Measurement of Asterion– Porion length (AST-PO)





Figure :5 Measurement of Porion– Mastoidale length (PO-MS)

## RESULTS

All the parameters measured, i.e., mastoid length, mastoid breadth, the porion-mastoidale, mastoidale-asterion, asterion-porion length, proved to have a higher value in males as compared to females and the differences were statistically significant for all these parameters

Following observations were made:

**Mastoid length:** was  $31.87 \pm 4.35$  mm in males and  $29.99 \pm 4.05$  mm in female.

**Mastoid breadth:** was  $22.57 \pm 4.10$  mm in males and  $21.54 \pm 3.55$  mm in females.

4 Asterion– Mastoidale length (AST-MS): was  $48.87 \pm 5.40$  mm in males and  $47.49 \pm 5.09$  mm in females.

Asterion– Porion length (AST-PO): was  $45.23 \pm 3.33$  mm in males and  $44.02 \pm 4.26$  mm in females.

4 Porion– Mastoidale length (PO-MS): was  $31.60 \pm 3.99$  mm in males and  $30.18 \pm 3.42$  mm in females.

**Table-1: Morphometric Parameters of the Mastoid Process of Male and Female Skulls**

Parameters (in mm)	Male (n=160)		Female (n=80)		p value
	Mean	± SD	Mean	± SD	
Mastoid Length	31.87	± 4.35	29.99	± 4.05	0.001
Mastoid Breadth	22.57	± 4.10	21.54	± 3.55	0.045
Asterion-Mastoidale	48.87	± 5.40	47.49	± 5.09	0.055
Asterion-Porion	45.23	± 3.33	44.02	± 4.26	0.017
Porion-Mastoidale	31.60	± 3.99	30.18	± 3.42	0.005

**Table-2: Morphometric Parameters of the Mastoid Process of both Sides of Male and Female Skulls**

Parameters (in mm)	Male (n=80)			Female (n=40)		
	Right	Left	p value	Right	Left	p value
	Mean±SD	Mean±SD		Mean±SD	Mean±SD	
<b>Mastoid Length</b>	32.07±4.53	31.67±4.18	0.563	30.31±4.41	29.68±3.99	0.854
<b>Mastoid Breadth</b>	22.74±4.00	22.42±4.22	0.625	22.07±3.79	21.01±3.26	0.183
<b>Asterion-Mastoidale</b>	48.99±4.98	48.76±5.83	0.795	47.62±5.19	47.37±5.06	0.826
<b>Asterion-Porion</b>	45.60±3.60	44.87±3.03	0.169	44.34±4.45	43.71±4.10	0.511
<b>Porion-Mastoidale</b>	32.14±4.02	31.08±3.91	0.093	30.66±3.52	29.71±3.31	0.219

## DISCUSSION

Analysis of the characteristics of the mastoid process is important in the determination of sex for forensic purposes and anthropologists. In the present study, mastoid length and breadth were more in males than females. When sex-wise analysis was done, the differences were found to be statistically significant but it was not found to be statistically significant on side-wise analysis. (Saadia et al 2016)

reported, the mean mastoid length was  $3.70 \pm 0.11$  cm in male and it was  $3.07 \pm 0.38$  cm in female. While (Passey et al 2015) and (Noack 2015) reported lower results in Asian races (mean mastoid length was 2.97 cm in male and 2.45 cm in female). (Saadia et al 2016) reported, the mean mastoid breadth was higher in male ( $2.80 \pm 0.24$  cm) than in female ( $2.31 \pm 0.29$  cm). (Nagaoka et al 2008) on Japanese skulls reported that the mean mastoid breadth was  $2.40 \pm 0.25$  cm in male and  $2.21 \pm 0.26$  cm in female. While (Sumati et al 2010) reported lower results on North Indian skulls, the mean of mastoid breadth was  $11.46 \pm 2.7$  mm in male and  $8.68 \pm 2.59$  mm in female.

The mean AST-MS length was  $48.87 \pm 5.40$  mm in males and  $47.49 \pm 5.09$  mm in females. It was more in males as compared to the females and showed statistically significant difference. On the left side it was  $48.76 \pm 5.83$  mm in male and  $47.37 \pm 5.06$  mm in female while on the right side it was  $48.99 \pm 4.98$  mm in male and  $47.62 \pm 5.19$  mm in female. In a study done by (Saadia et al 2016) the mean AST-MS length was  $5.06 \pm 0.28$  cm on left side and  $5.22 \pm 0.31$  cm on the right side in male and it was  $4.39 \pm 0.29$  cm on left side and  $4.44 \pm 0.35$  cm on right side in female. (Jain et al 2013) on Indian skulls reported that the mean AST-MS length was higher in male ( $4.92 \pm 0.80$ ) than in female ( $4.47 \pm 0.72$ ) on both sides. But (Suazo et al 2008) found that the mean AST-MS length was nearly similar in male ( $5.02 \pm 0.49$ ) and female ( $5.01 \pm 0.51$ ) in Brazilian skulls. The mean AST-PO length was  $45.23 \pm 3.33$  mm in males and  $44.02 \pm 4.26$  mm in females. It was more in males as compared to the females and statistically significant difference was observed. On the left side it was  $44.87 \pm 3.03$  mm in male and  $43.71 \pm 4.10$  mm in female while on

the right side it was  $45.60 \pm 3.60$  mm in male and  $44.34 \pm 4.45$  mm in female. (Saadia et al2016) reported mean asterion- porion (AST-PO) length was higher in male ( $4.66 \pm 0.32$  cm on left side and  $4.56 \pm 0.22$  cm on right side) than in female ( $4.26 \pm 0.21$  cm on left side and  $4.23 \pm 0.19$  cm on right side) on both sides. In another study done by (Jaja et al 2013) on Nigerian skulls in which there was significant difference between male ( $4.60 \pm 0.71$  cm) and female ( $4.30 \pm 0.65$  cm) in mean of AST-PO length on left side but in present study it was slightly higher in males than females. (Bhaskar et al2013) found that the mean mastoid length was  $3.56 \pm 0.39$  cm in male and  $3.05 \pm 0.40$  cm in female in South Indian skulls. The mean PO-MS length was  $31.60 \pm 3.99$  mm in males and  $30.18 \pm 3.42$  mm in females. On the left side it was  $31.87 \pm 3.91$  mm in male and  $29.71 \pm 3.31$  mm in female. On the right side it was  $32.14 \pm 4.02$  mm in male and  $30.66 \pm 3.51$  cm in female. (Saadia et al2016) reported mean PO-MS length on the left side was  $3.25 \pm 0.12$  cm in male and  $2.63 \pm 0.27$  cm in female while on the right side it was  $3.29 \pm 0.14$  cm in male and  $2.76 \pm 0.27$  cm in female. In the present study it was more in males as compared to the females and showed statistically significant difference. On comparing with the results of other studies, the present study shows that the parameters of the mastoid process measured can be accountable in medico-legal investigations, and it can be taken as a sex indicator among North Indians.

## CONCLUSION

The study on the parameters of the mastoid process is important in the determination of sex for forensic purposes and anthropologists. It was concluded in our study that the mean mastoid parameters were more in male skulls than female skulls.

FUNDING; None



## REFERENCES

1. Bhaskar B., Nidugala H, Avadhani R(2013). Mastoid process –Atool for sex determination, an anatomical study in South Indian skulls. *IJBR* 4(2)106-110.
2. Jaja B.N., Ajuja C.O. and Didia B.C.( 2013) Mastoid triangle for sex determination in adult Nigerian Population: a validation study. *JFS* 58(6):1575-8.
3. Noack T. B.( 2015) Sexual dimorphism in the crania in a Norwegian sample. B.A, University of Texas at San Antonio. p.24-28.
4. Nagaoka T., Shizushima A., Sawada J., Tomo S., Hoshino K. and Sato H.( 2008) Sex determination using mastoid process measurements: standards for Japanese human skeletons of the medieval and early modern periods. *Anthropological Sci.* 116:105–113.
5. Passey J, Mishra SR, Singh R, Sushobhana K, Singh S, Sinha P(2015). Sex determination using mastoid process. *AJMS*;6(6):93-5.
6. Saadia A. Shalaby, Essam M, Eid, Omar A. (2016) Morphometric Study Of Mastoid Canal And Suprameatal Triangle Of Human Egyptian Skull, With Gender Determination. *Nature and Science.* 14 (4): 67-73.
7. Standring S.(2005) Skull and Mandible In: Standring S(ed.) Gray's Anatomy. The Anatomical basis of clinical practice. 39<sup>th</sup> edition; Edinburg; Elsevier,. p. 470.
8. Suazo G.I.C., Zavando M.D.A. and Smith R.L.(2008): Sex determination using mastoid process measurements in Brazilian skulls. *Int J Morphol.* 26:941–944
9. Sumati, Patnaik VVG, and Phatak A.( 2010) Determination of sex from mastoid process by discriminant function analysis. *JASI*;59(2):222-28.

