

# **A STUDY OF CLINICAL FEATURES EVALUATION AND SURGICAL MANAGEMENT OF VARICOSE VEINS OF LOWER LIMBS**

## **ABSTRACT**

**BACKGROUND:** Varicose veins have been described as chronic venous disorder since ancient times. The number of patients coming to hospital for treatment is far less than the actual incidence as they come to hospitals only when complications arise

**OBJECTIVES:** To study the clinical presentations, surgical management and its outcome and complications associated with varicose veins in lower limbs.

**METHODS:** A total of 50 patients with primary varicose veins were sourced from cases admitted to Basaveshwar Teaching and General Hospital attached to Mahadevappa Rampure Medical College, Gulbarga from 1<sup>st</sup> August 2022 to 31<sup>st</sup> January 2024. They were investigated, operated and followed up, and the outcome evaluated. All the information was taken down in the proforma designed for the study.

**RESULTS:** The study observed that majority patients 44% belonged to age group of 41-60 years, Male to Female ratio being 7.3:1. The most common presenting symptom being pain 86% followed by edema 52%, most common usg finding being perforator incompetence 92%. In the study 94% patients underwent Subfascial Perforator Ligation, 50% underwent only Saphenofemoral flush ligation. Study revealed that 18% patients had complications of wound infection, thrombophlebitis, hematoma in the immediate post operative period. In follow up after 3 months 6% patients had Saphenous Neuralgia.

**CONCLUSION:** The prevalence of varicose veins is most common in middle aged individuals with males being more commonly affected than females, prolonged standing was observed as the significant risk factor. Surgical interventions like the Trendelenburg operation with Stripping and Subfascial Perforator ligation are highly effective for treating varicose veins,

29 Bisgaard's method is effective for Healing Ulcers. Most postoperative complications were  
30 managed conservatively and showed excellent resolution with time. No patient developed life  
31 threatening or debilitating complications and there was no mortality or recurrence in this study.

32 **KEYWORDS:** Varicose Vein, Duplex, Venous Ulcer, Sapheno-femoral, Perforators,  
33 Sapheno-popliteal, Incompetence, Flush Ligation, Saphenous Neuritis,  
34 Complications.

## 35 INTRODUCTION

36 Varicose veins and their associated symptoms and complications constitute the most common  
37 chronic vascular disorders leading to surgical treatment. —varicosity is the penalty for  
38 verticality against gravity<sup>1</sup> . "Varicose" is derived from the Latin word for "dilated." d."  
39 Definitions of varicose veins vary considerably: Arnoldi describes them as "clearly visible,  
40 dilated, tortuous, and possibly prominent subcutaneous veins of the lower extremities," while  
41 Dodd and Cockett refer to them as "veins with a loss of valvular efficiency." The World Health  
42 Organization (WHO) defines them as "veins with a saccular dilatation, often tortuous.

## 43 EPIDEMIOLOGY

44 Varicose veins are common. The prevalence has been variously reported from as little as 2% to  
45 over 20% in population studies<sup>2</sup>

### 46 Objectives of Study

- 47 1. To study age and sex incidence of varicose veins of lower limbs and their clinical features
- 48 2. To study the correlation between occupation and incidence of varicose veins in lower limb
- 49 3. To study different surgical methods like Sapheno -femoral flush ligation, multiple subfascial
- 50 perforator ligation, stripping of veins and respective clinical outcomes at the end of three months
- 51 of follow up period.

## 52 MATERIALS AND METHODS

53 The present clinical study and management of varicose veins was done at Mahadevappa  
54 Rampure Medical College Gulbarga during period of AUG 2022-JAN 2024

Source of Data
During the study period 50 inpatients with Varicose veins admitted in Department of

General Surgery, Basaveshwar Teaching & General Hospital attached to Mahadevappa Rampure Medical College, Kalaburagi.were studied .

### **Methods of collection of data**

**Study Design :** Prospective Interventional Study

**Study Settings :** Department of General Surgery, Basaveshwar Teaching and General Hospital, Kalaburagi

**Sample Size : 50**

**Sampling Procedure :**

p=incidence of Varicose veins is 15- 20%

q= 100-p=100-15=85

L= permissible error of p

L=50% of p=10

Sample size  $n = (Z^2 \alpha pq) / L^2$

$$= [(1.96)^2 \times 15 \times 85] / 10^2$$

$$= 48.98$$

$$= 50$$

**Duration of Study :** (18 months), from 1<sup>st</sup> AUGUST 2022 –31<sup>st</sup> JANUARY 2024

**Inclusion Criteria :**

- Patients presenting with symptoms such as aching heaviness cramps
- Patients with complications of Dermatitis Ulceration superficial thrombophlebitis
- Patients with cosmetic concerns

**Exclusion Criteria :**

- All patients with secondary varicose veins with Deep Vein Thrombosis ,Pregnancy, Peripheral Arterial diseases
- All patients with recurrent varicose veins

**Informed Consent**

Written and informed consent was taken in patients own vernacular language

**.Statistical Analysis**


Statistical analysis was done using IBM SPSS software version 20.0 Chi square test was applied for qualitative analysis. Student t test and ANOVA was applied for quantitative data analysis. If  $p < 0.05$  was considered significant.

55 The routine investigations were done and special investigations were performed wherever  
56 necessary. The pre-operative treatment, operative findings and post-operative outcome are  
57 documented. Routine follow up was done during the immediate post-operative period and every  
58 day till discharge.

**RESULTS****Table No.1: Age wise distribution of patients**

Age in years	Number of patients	Percentage
≤ 20	2	4.0
21—40	19	38.0
41—60	22	44.0
61—80	7	14.0
<b>Total</b>	50	100.0
<b>Mean ± SD</b>	43.66 ± 13.52	

62 Study observes that, majority of patients each 22 (44.0%) belong to the age group of 41—60  
63 years. Followed by 19 (38.0%) of patients belong to the age group of 21—40 years, 7 (14.0%) of  
64 patients belong to the age group of 61—80 years and 2 (4.0%) of patients age is in the range was  
65  $\leq 20$  years. Minimum age of patient was 18 years and maximum age of patient was 70 years. The  
66 mean age of patients was 43.66 years




67

68 **Graph 1: Pie chart represents gender wise distribution of patients**

69 Study observed that; Male patients were predominant in the study they were 44 (88.0%) and 6  
70 (12.0%) of patients were females. Male to Female ratio was 7.3:1

71



72

73

**Graph 2: Pie diagram presents occupation wise distribution of patients**

74

According to occupational status, majority of the cases were seen in the prolonged standing workers. In this study, manual labourers were found constituting 10 (20.0%) of cases, shopkeepers of business 7 (14.0%), employees 12 (24.0%), and farmers or agriculturalist 5 (10.0%)

75

76

77


78

**Table No2 : Distribution of patients based on side of incidence of varicosity**

Side	Number of patients	Percentage
Right	26	52.0
Left	21	42.0
Bilateral	3	6.0
Total	50	100.0

79

80 Study observed that; an increased incidence of varicosity was noted on right side i-e 26 (52.0%),  
 81 the incident of varicosity were seen on left side 21 (48.0%) and 3 (6.0%) of patients were seen  
 82 the bilateral incidence of varicosity



83  
 84 **Graph3:Bar diagram represent symptoms wise distribution of varicosity patients**

85 The most common presenting symptom in varicose veins in this study was pain seen in 42  
 86 (82.0%) of cases. Oedema was reported in 26 (52.0%) cases. In this study, 13 (26.0%) cases  
 87 presented with pigment and ulcer was seen in 5 cases (10.0%).

88  
 89 **Table No.3: USG finding wise distribution of varicosity patients**

90

USG Findings	Number of patients	Percentage

<b>SFI</b>	<b>Present</b>	24	48.0
	<b>Absent</b>	26	52.0
<b>PI</b>	<b>Present</b>	46	92.0
	<b>Absent</b>	4	8.0
<b>SPI</b>	<b>Present</b>	4	8.0
	<b>Absent</b>	46	92.0

91 The most common USG findings observed were Perforator incompetence in 46 (92.0%) of  
 92 cases, followed by saphenofemoral junction incompetence were observed in 24 (48.0%) of  
 93 cases and saphenopopliteal junction incompetence were observed in 4 (8.0%) of cases.

94

95 **Table No.4: Treatment procedure wise distribution of varicosity patients**

<b>Treatment procedure</b>	<b>Number of patients</b>	<b>Percentage</b>
<b>Bisgaard</b>	5	10.0
<b>Saphenofemoral flush ligation</b>	25	50.0
<b>Stripping of veins</b>	14	28.0
<b>Subfascial perforator ligation</b>	47	94.0
<b>Saphenopopliteal junction ligation</b>	4	8.0

96

97

**Table 5: Surgical procedures performed**


<b>Surgical procedures</b>	<b>Limb</b>	<b>Percentage</b>
Sapheno-femoral flush ligation + Stripping	4	8
Sapheno-femoral flush ligation + Stripping + Subfascial	7	14



ligation		
Sapheno-femoral flush ligation + Subfascial ligation of perforators	8	16
Sapheno-femoral flush ligation + Stripping + sapheno popliteal ligation+ Subfascial ligation	2	4
Subfascial ligation of perforators	23	46

98

99 In this study 47 (94.0%) of patients underwent the surgical procedure of subfascial perforator  
 100 ligation, followed by 25 (50.0%) of patients underwent Sapheno-femoral flush ligation. 14  
 101 (28.0%) of patients underwent stripping of veins. 5 (10.0%) of patients underwent Bisgaard  
 102 regime for venous ulcer before surgical procedure and 4 (8.0%) of patients underwent  
 103 Sapheno-popliteal junction ligation.



104

105 **Graph 4:Bar diagram presents venous system involved distribution of patients**

106

**Table No.6: Perforator type involved in varicose vein patients**

Perforator type	Number of patients	Percentage
BK, Ankle	23	46.0
Multiple	11	22.0
BK	10	20.0
Low leg and Ankle	2	4.0

107  
108 Study reveals that; majority of patient's 23 (46.0%) perforator type was below knee and ankle  
109 perforator incompetence . Followed by 11 (22.0%) of patients had Multiple perforator  
110 incompetence, 10 (20.0%) of patients had only below knee perforator incompetence, 2 (4.0%)  
111 of patients had only ankle perforator incompetence and 4 (8.0%) of patients did not have  
112 perforator incompetence

113 **Table No.7: Distribution of varicose vein patients based on early complications**

Early complications	Number of patients	Percentage
Wound Infection	4	8.0
Thrombophlebitis	3	6.0
Hematoma	2	4.0
Total	9	18.0

114  
115 Study observed that; 9 (18.0%) of patients had seen complications. 4 (8.0%) of patients had  
116 wound infection, 3 (6.0%) of patients had thrombophlebitis and 2 (4.0%) of patients had  
117 hematoma.

118 **DISCUSSION**

119 **AGE DISTRIBUTION**

Studies	Age range(yr)
Present study	18-70
Malhotra et al <sup>3</sup>	18-65

120 Table 8; comparison study for agr distribution

121 Malhotra et al<sup>3</sup> (1972) in their study comprising 677 patients from both North and South India  
122 had an age range of 18-65  
123 years. In the West Wright et al<sup>4in</sup> their study of 1338 patients in England had an age range of 20-  
124 75 years. This is also corroborated in the studies by Lateef et al (65%), Fegan et al (75%) and  
125 Ratkal et al (72%). Maximum incidence being 41-60 yrs age group with mean age of patients  
126 being 43.66 yrs . So it affects the bread-earning members of the family, causing socioeconomic  
127 problems. This also indicates earlier health- seeking behavior in the present generation.

128 **SEX DISTRIBUTION: Table 9:comparison study for sex distribution**

Studies	Male: Female
Present study	7.3:1
Widmer et al <sup>5</sup>	1:1
Callam et al <sup>6</sup>	1:2

129 In my series male to female ratio was found to be 7.3:1. Widmer<sup>5</sup> in Switzerland recorded a  
130 ration of 1:1. Callam et al<sup>6</sup> in England recorded a ratio of 1:2. . It may be because, they do not  
131 undergo the occupational hazards of that of males, like prolonged standing, physical stress  
132 involving increased intraabdominal pressure. The decreased occurrence of disease in females at  
133 our set up may be because females may be resistant to complications of varicose veins probably  
134 due to less average height compared to male which has a direct impact on venous hypertension .

135 **Table 10:Comparison study of occupation versus varicose veins**

Occupation	Lateef (1971) <sup>8</sup>	Ratkal (1980) <sup>9</sup>	Present series
------------	----------------------------	----------------------------	----------------

Occupation involving prolonged standing	35%	44%	52%
---	-----	-----	-----

136 Varicose veins are more common in persons, whose occupation force them to stand for  
 137 prolonged hours. In the present study about 52% of patients had occupations, which involved  
 138 prolonged standing like farmer, policemen, bus driver etc

139 **Table 11 :Comparison of incidence of Clinical Features**

Symptoms	Present series		Rudofsky G. Langenbecks Arch Chir (%) <sup>10</sup>	O' Shaughnessy M et. Al (%) <sup>11</sup>
	No of patients	% C		
Prominent veins	50	100	90	92
Prominent veins and pain	42	84	30	54
Prominent veins and edema	26	52	52	62
Pigmentation	13	26	13	22
Venous ulceration	5	10	9	14
Previous history of DVT	-	-	-	5

140

LIMB	A. H. M. Dur, A. J. C. Mackaay <sup>65</sup> et al <sup>7</sup>	PRESENT STUDY
RIGHT	48.55%	52%
LEFT	51.45%	42%

141

**Table 12:comparison study of limb involvemen**

142 **SITE OF PERFORATOR INCOMPETENCE:**

143 Around 22% of the patients had multiple perforator incompetence. Patients who had multiple  
 144 perforator incompetence had one or the other complications of varicose veins. Isolated below  
 145 knee perforator incompetence was seen in 20%, The results observed were similar to study by  
 146 Labropoulos et al

147

**Table 13: Comparison of Perforator Incompetence:**

Studies	Incompetent perforator(%)
Present study	92%
Labropoulos N et al <sup>12</sup>	68%

148

**Table 14: Complications post surgery according to**

149

**Hagmuller G.W and Langenbecks Archchis study<sup>13</sup>**

Complications	Percentage	Present study
---------------	------------	---------------

Femoral vein injury	1%	0%
Femoral artery injury	0.02%	0%
Deep vein thrombosis	0.15%	0%
Pulmonary embolism	0.06%	0%

150

151 In the present study some minor complications occurred which were managed conservatively.

152 The study conducted by Hagmuller G.M. showed incidence of some major complications which

153 are very rare and none of which occurred in the present study group.

154 **FOLLOW UP :**

155 At 3 month, 6% of the patients had complications. The complications looked for were persistent

156 oozing, stitch abscess, neuralgia, thrombophlebitis and recurrence. Neuralgia was seen in 4%

157 patients and was treated with cobalamin capsules. Patients were advised elastic compression

158 stockings for 1 year post operatively. None of the patients developed recurrence.

## 159 **CONCLUSION**

160 This detailed study of 50 cases of lower limb varicose veins led to the following conclusions:

161 • Varicose veins of the lower limb are a fairly common condition, though the number of  
162 cases reported to hospitals is likely lower than the true incidence, as many patients do not  
163 seek treatment unless symptoms are present.

164 • The most commonly affected age group is between 41 and 60 years. Most patients  
165 presented to the hospital due to complications rather than for cosmetic reasons.

166 • The majority of patients were male. The lower number of females in the study could be  
167 due to cultural factors, such as women being less concerned about cosmetic appearance  
168 or potentially having a lower risk of complications due to hormonal differences, shorter  
169 average height, or less intense physical activity.

170 • While no definitive conclusions about etiology can be drawn from this small sample,  
171 there is a clear relationship between occupation and the incidence of varicose veins.

172 • The long saphenous vein is more commonly affected than the short saphenous vein. Sub-  
173 fascial ligation of incompetent perforators yields satisfactory results.

174 • Clinical examination has a high predictive accuracy and can provide sufficient  
175 information for treatment, especially in centers where color Doppler imaging is not  
176 available or affordable.

- 177 • The use of color Doppler is a valuable supplement to clinical examination and is strongly  
178 recommended to prevent recurrences and reduce morbidity, as it effectively detects  
179 venous incompetence.
- 180 • The saphenopopliteal junction (SPJ) is highly variable and should always be marked  
181 preoperatively using Doppler.
- 182 • The outcomes of cases of primary varicose veins depend on thorough and complete  
183 clinical examination and duplex scanning by an experienced radiologist.
- 184 • In the absence of junctional incompetence, sub-fascial ligation of incompetent perforators  
185 was associated with no recurrences.
- 186 • Complications are minimal when cases are meticulously selected and operated on,  
187 enabling patients to lead a near-normal life post-surgery, with a very low mortality rate.
- 188 • While newer treatment methods for varicose veins show promising results, they require  
189 long-term follow-up and may not be affordable for many patients due to cost factors.

## 190 **BIBLIOGRAPHY**

- 191
- 192 • Russell RCG, Williams NS, Bulstrode CJK, —Venous disorders| in Bailey and Love’s  
193 Short practice of surgery, Ch.54; 28th Edn; Arnold publications; 2023: 925-943.
- 194 • Patrick H carpentier —Prevalence, risk factors, and clinical patterns of chronic venous  
195 disorders of lower limbs: A population-based study in France| Journal of  
196 Vascular Surgery.2004;40:650-659.
- 197 3 S L Malhotra —An Epidemiological Study of Varicose Veins in Indian Railroad  
198 Workers from the South and North of India, with Special Reference to the  
199 Causation and Prevention of Varicose Veins|. International J. Of Epidemiology  
200 1972; (1): 177-183.
- 201 4 Wright et al. —The prevalence of venous disease in a west London population. In:  
202 Davy A, stemmer R, Eds. Phlebology‘ 89. Paris : libbey Eurotext, 1989: 176-8.

- 203 5 Widmer LK ed. Peripheral venous disorders prevalence and socio-medical  
204 importance. Bern: Hans Huber, 1978:1-90.
- 205 6 Callam, M. J., —Epidemiology of varicose veins. British Journal of Surgery,  
206 1994: 81; 167–173.
- 207 7 A.H.M. Dur, A.J.C Mackay et al. —Duplex assessment of clinically diagnosed  
208 chronic venous insufficiency. Br. J. surg. June 1992; 79: S, 155-161.
- 209 8 Lateef MA. Clinical pathological study of the primary varicose veins in the lower limb,  
210 Br.J.Surg. 1995; 82: 855-856.
- 211 9 Ratkal C. Thesis on clinical Pathological study of varicose veins of the lower limb. University  
212 of Mysore, 1980.
- 213 10 Rudofsky G. Epidemiology and pathophysiology of primary varicose veins. Langenbecks  
214 Arch Chir 1988; Suppl 2:139-44
- 215 11 O' Shaughnessy et al. —Surgery in the treatment of varicose veins. Ir.Med.J,1989 June;  
216 82(2): 54-55.
- 217 12 Labropoulos N. et al. —Where does venous reflux start J Vasc surg 1997; 26(S) 738- 742.
- 218 13 Hagmuller GW. Complications in surgery of varicose veins. Langenbecks  
219 Arch Chir Suppl Kongressbd 1992; 470-4.

220

221

222

223

224

225

226

227